

HIGHER DIMENSIONAL ALGEBROIDS AND
CROSSED COMPLEXES

Thesis submitted to the University of Wales in support of
the application for the Degree of Philosophiae Doctor

By

GHAFAR HUSEEN MOSA

Supervised by Professor

R. Brown

1980 MATHEMATICS SUBJECT CLASSIFICATION: 18D05 , 18E05 ,
55U99 .

KEY WORDS: Algebroid , Double Algebroid , ω -Algebroid ,
Crossed module , Crossed complex .

G.H.Mosa

Department of Pure Mathematics ,
University College of North Wales ,
Bangor ,

Gwynedd LL57 2UW

U.K.

November , 1986

DECLARATION

The work of this thesis has been carried out by the candidate and contains the results of his own investigations . The work has not already been accepted in substance for any degree , and is not being concurrently submitted in candidature for any degree . All sources of information have been acknowledged in the text .

R. Brown

DIRECTOR OF STUDIES

Mora

CANDIDATE

ACKNOWLEDGEMENT

I would like to thank my supervisor , Professor R.Brown for suggesting the problems from which this thesis arose , and for his help and encouragement during the course of this work . Thanks are also due to Dr. T.Porter for his help and helpful discussions .

I am grateful to the Iraqi goverment for providing me with a maintenance grant . Finally , I am grateful for my parents for their help .

G.H.MOSA

SUMMARY

The equivalence between the category of crossed modules (over groups) and the category of special double groupoids with connections and with one vertex was proved by R.Brown and C.B.Spencer . More generally , C.B.Spencer and Y.L.Wong have shown that there exists an equivalence between the category of 2-categories and the category of double categories with connections .

R.Brown and P.J.Higgins have generalised the first result : they proved that there exists an equivalence between the category of ω -groupoids and the category of crossed complexes (over groupoids) .

In this thesis we develop a parallel theory in a more algebraic context , with expectation of applications in non-abelian homological and homotopical algebra . We prove an equivalence between the category of crossed modules (over algebroids) and the category of special double algebroids with connections . Moreover we prove a similar result for the 3-dimensional case , that is , we prove that there exists an equivalence between the category $(\text{Crs})^3$ of 3-truncated crossed complexes and the category $(\omega\text{-Alg})^3$ of 3-tuple algebroids . Also we end this work by giving a conjecture for the higher dimensional case . In particular , we have

Theorem: The functors γ , λ form an adjoint equivalence

$$\gamma : \underline{DA}^! \longleftrightarrow \underline{C} : \lambda$$

where $\underline{DA}^!$ is the category of special double algebroids with connections and \underline{C} is the category of crossed modules .

Theorem: The functors γ , λ form an adjoint equivalence

$$\gamma : (\omega\text{-Alg})^n \longleftrightarrow (\text{Crs})^n : \lambda$$

for $n = 3 , 4$.

Finally we give a conjecture whose validity would be sufficient for the general equivalence of categories of ω -algebroids and crossed complexes .

In chapter VI we explain some results which have been obtained in the case of groupoids and higher dimensional groupoids , and suggest the possibility of obtaining similar results in the case of algebroids and higher dimensional algebroids .

CONTENTS

INTRODUCTION

CHAPTER I : R-ALGEBROIDS

\$0. INTRODUCTION	1
\$1. R-ALGEBROIDS	3
\$2. CROSSED MODULES (OVER ASSOCIATIVE ALGEBRAS) ...	8
\$3. CROSSED MODULES (OVER ALGEBROIDS)	10

CHAPTER II : DOUBLE R-ALGEBROIDS

\$0. INTRODUCTION	15
\$1. DEFINITIONS	15
\$2. FUNCTORS (DOUBLE ALGEBROIDS) \rightarrow (CROSSED MODULES) ...	23
\$3. EXAMPLES	26

CHAPTER III : THE EQUIVALENCE BETWEEN THE CATEGORY C OF CROSSED MODULES AND THE CATEGORY DA! OF SPECIAL DOUBLE ALGEBROIDS WITH CONNECTIONS

\$0. INTRODUCTION	30
\$1. THIN STRUCTURES AND CONNECTIONS	31
\$2. THE FOLDING OPERATION	41
\$3. THE FUNCTOR $\lambda : \underline{C} \dashrightarrow \underline{DA}!$	46
\$4. THE EQUIVALENCE OF CATEGORIES	52
\$5. REFLECTION	56

CHAPTER IV : ω -ALGEBROIDS (WITHOUT CONNECTIONS) AND CROSSED COMPLEXES

\$0. INTRODUCTION	60
\$1. ω -ALGEBROIDS (WITHOUT CONNECTIONS)	60
\$2. CROSSED COMPLEXES	63
\$3. THE FUNCTOR $(\omega\text{-Alg}) \dashrightarrow (\text{Crs})$	66

**CHAPTER V : THE EQUIVALENCE BETWEEN n -TUPLE ALGEBROIDS
AND CROSSED COMPLEXES FOR $n=3$ and 4**

\$0. INTRODUCTION	72
\$1. ω -ALGEBROIDS WITH CONNECTIONS	72
\$2. FOLDING OPERATIONS	75
\$3. COSKELETON OF ω -ALGEBROIDS	98
\$4. THE EQUIVALENCE OF CATEGORIES	102

**CHAPTER VI : CONJECTURED RESULTS ON ALGEBROIDS ,
 ω -ALGEBROIDS AND CROSSED COMPLEXES 105**

APPENDIX I	108
APPENDIX II	112
APPENDIX III	118
APPENDIX IV	120
APPENDIX V	125
APPENDIX VI	128
REFERENCES	134

INTRODUCTION

A. AIMS AND BACKGROUND:

1. Overall aim:

There are many useful analogies between the theory of groups and the theory of algebras which are exploited for example in homological algebra . Some interesting generalisations of groups are groupoids , crossed modules , crossed complexes , double groupoids and ω -groupoids , dating respectively from 1926 [Brandt - 1] , 1946 [J.H.C.Whitehead - 1,2] , 1949 [Blakers - 1] , 1965 [Ehresmann - 1] and 1977 [Brown-Higgins - 8] . Corresponding to groupoids as generalisations of groups , we have algebroids as generalisations of algebras , a theory due to B.Mitchell (1972) . There are also notions of crossed modules of algebras . But a theory of double and n -tuple algebroids does not seem to be available , and it is our aim to investigate this idea .

In order to see the motivation for this investigation and the kind of result to be expected , we first recall some facts on the group case .

2. Crossed modules , crossed complexes in groups and ω -groupoids:

First , a group homomorphism $\partial: M \rightarrow P$ is said to be a crossed P -module (in groups) if there is given an action of P on M , $(p,m) \mapsto Pm$ which satisfies the following axioms :

(i) $\partial(Pm) = p (\partial m) p^{-1}$ (ii) $\partial mm' = m m' m^{-1}$ for $m,m' \in M$
and $p \in P$. Standard examples of crossed modules are :

- 1) the inclusion $N \rightarrow P$ of a normal subgroup N of the group P , with the action of P on N given by conjugation ;
- 2) the zero morphism $0 : M \rightarrow P$ in which M is a P -module in the usual sense ;
- 3) the boundary map $\partial : \pi_2(X, Y, x_0) \rightarrow \pi_1(Y, x_0)$ from the second relative homotopy group to the fundamental group with the standard action of $\pi_1(Y, x_0)$ on $\pi_2(X, Y, x_0)$.

As this last example suggests , crossed modules can be used to model certain homotopy types . In fact from the standpoint of homotopy theory , crossed modules should be viewed as "2-dimensional groups" . It is reasonable to ask then , what are the n -dimensional groups (or crossed modules) ?

J.H.C.Whitehead gave a partial answer to this by introducing what he called a "homotopy system" , but which are now called crossed complexes . These gadgets consist of a sequence of groups

$$\dots \xrightarrow{\partial_n} C_n \xrightarrow{\partial_{n-1}} C_{n-1} \xrightarrow{\partial_{n-2}} \dots \xrightarrow{\partial_3} C_3 \xrightarrow{\partial_2} C_2 \xrightarrow{\partial_1} C_1 \xrightarrow{\partial_0} C_0$$

where C_0 is a single point and satisfy the axioms ;

- i) ∂_1 is a crossed module ;
- ii) C_n is abelian for $n \geq 3$;
- iii) $\partial^2 = 0$;
- iv) C_1 acts on C_n , $n \geq 2$ and $\partial_1 C_2$ acts trivially on C_n for $n \geq 3$.

The standard example of a crossed complex is obtained from a pointed filtered space (c.f. [Br-3]) .


Work in homotopy theory has developed the well known notion of "groupoids" , which are categories in which every arrow is invertible .

Since a crossed module has been considered as a "higher dimensional group" , the question arises : what is a higher dimensional groupoid ? Ehresmann [Eh-1] has defined the notion of double groupoid . R-Brown and C.B.Spencer have proved that there exists an equivalence between the category of crossed modules (over groups) and the category of double groupoids with special connections and one vertex . But the general case has been defined in [B-Hi-8] ; namely they have defined ω -groupoids and crossed complexes (over groupoids) by using the cubical set notion . Moreover they have proved in [B-Hi-2] there exist an equivalence between the category of ω -groupoids and the category of crossed complexes (over groupoids) .

The above discussion of the development of group theory in this direction is summarised in the diagram

Groups \rightarrow Groupoids \rightarrow Double groupoids \rightarrow ω -groupoids .

There are in fact a remarkable collection of equationally defined categories of (many-sorted) algebras which are non-trivially equivalent to ω -groupoids . These are summarised in the following diagram :



in which the arrows denote explicit functors which are equivalence of categories . The symbols in square brackets give references to the proofs .

3. Crossed modules and Crossed complexes over algebras:

The work of [Ge-1] essentially involves the notion of crossed modules in associative and commutative algebras under a different name . Also the work of [K-L-1] in algebraic K-theory has introduced crossed modules of Lie algebras . The definition of crossed modules in associative algebras is given on page (9,10) .

The notion of crossed modules of algebras has been generalised to crossed complexes over algebras [Po-3] , namely ;

Let R be a commutative ring and let K be an R -algebra . A crossed complex of R -algebras is a sequence of R -algebras

$$\dots \rightarrow C_n \xrightarrow{\partial_n} C_{n-1} \rightarrow \dots \xrightarrow{\partial_2} C_2 \xrightarrow{\partial_1} K$$

in which

- i) ∂_1 is a crossed K -module ,
- ii) C_i for $i \geq 1$ is a K -module on which $\partial_1 C_1$ operates trivially
- iii) for $i \geq 1$, $\partial_{i+1} \partial_i = 0$.

Now one can ask , what are the higher dimensional algebras ? In this thesis we shall give a partial answer to this question and we will give some extra conjectures .

4. Algebraic geometry:

The idea of this work arose from the consideration of bringing crossed module ideas into commutative algebra and algebraic geometry ; namely an ideal in a polynomial ring corresponds to an affine algebraic variety . Crossed modules in commutative algebras are generalisations of ideals .

One would like to know the geometric analogue of a crossed module , but nothing seems to be known on this question .

The original motivation for this thesis was to see if it would be easier to find analogues of "double commutative algebroids" in algebraic geometry, assuming such were equivalent to crossed modules. This led to the problem of finding analogues for algebroids of the work of Brown - Higgins on ω -groupoids, and this problem has since occupied our full attention.

There are still many problems in relating this work to algebraic geometry, but we believe this will eventually be possible.

B. STRUCTURE AND MAIN RESULTS:

In chapter I we give an example to show how algebras are appropriately generalised to algebroids and we show that the category of R -algebroids is a monoidal closed category. We give the definition of a crossed module over an associative algebra and introduce the definition of a crossed module over an algebroid. Also we deduce some properties of crossed modules similar to the well known properties of crossed modules over groups.

In chapter II, we define an algebroid in one higher dimension. In fact we introduce the notion of a double algebroid by using double categories; namely a double R -algebroid D is four related R -algebroids

$$(D, D_1, \partial_1^i, c_1, +_1, *_1, \dots) , (D, D_2, \partial_2^i, c_2, +_2, *_2, \dots)$$

$$(D_1, D_0, \partial_1^i, c, +, *, \dots) , (D_2, D_0, \partial_2^i, c, +, *, \dots)$$

where $i = 0, 1$ and these algebroids satisfies the following

axioms :

$$i) \quad s_2^i \partial_2^j = s_1^j \partial_1^i \quad i, j \in \{0, 1\}$$

$$ii) \quad \partial_2^i(\alpha +_1 \beta) = \partial_2^i \alpha + \partial_2^i \beta, \quad \partial_1^i(\alpha +_2 \beta) = \partial_1^i \alpha + \partial_1^i \beta$$

$$\partial_2^i(\alpha *_1 \beta) = \partial_2^i \alpha *_1 \partial_2^i \beta, \quad \partial_1^i(\alpha *_2 \beta) = \partial_1^i \alpha *_2 \partial_1^i \beta$$

for $i = 0, 1$, $\alpha, \beta \in D$ and both sides are defined .

$$iii) \quad r \cdot_1 (\alpha +_2 \beta) = (r \cdot_1 \alpha) +_2 (r \cdot_1 \beta),$$

$$r \cdot_2 (\alpha +_1 \beta) = (r \cdot_2 \alpha) +_1 (r \cdot_2 \beta)$$

$$r \cdot_1 (\alpha *_2 \beta) = (r \cdot_1 \alpha) *_2 (r \cdot_1 \beta)$$

$$r \cdot_2 (\alpha *_1 \beta) = (r \cdot_2 \alpha) *_1 (r \cdot_2 \beta)$$

$$r \cdot_1 (s \cdot_2 \alpha) = s \cdot_2 (r \cdot_1 \alpha)$$

for $\alpha, \beta \in D$, $r, s \in R$ and both sides are defined .

$$iv) \quad (\alpha +_1 \beta) +_2 (\gamma +_1 \delta) = (\alpha +_2 \gamma) +_1 (\beta +_2 \delta)$$

$$(\alpha *_1 \beta) *_2 (\gamma *_1 \delta) = (\alpha *_2 \gamma) *_1 (\beta *_2 \delta)$$

$$(\alpha +_1 \beta) *_2 (\gamma +_1 \delta) = (\alpha *_2 \gamma) +_1 (\beta *_2 \delta)$$

$$(\alpha +_2 \beta) *_1 (\gamma +_2 \delta) = (\alpha *_1 \gamma) +_2 (\beta *_1 \delta)$$

for $\alpha, \beta, \gamma, \delta \in D$ and both sides are defined .

$$v) \quad c_1(a + a_1) = c_1 a +_2 c_1 a_1, \quad c_2(b + b_1) = c_2 b +_1 c_2 b_1$$

for $a, a_1 \in D_1$, $b, b_1 \in D_2$ and the additions are defined .

Thus we get a category of double R-algebroids DA .

We can ask now what is the relation between the category of crossed modules (over algebroids) and the category of double R-algebroids . At this stage we prove the following ;

Proposition: If D is a double R-algebroid , then we have two crossed modules associated with D . That is , there exist two functors from the category of double R-algebroids to the category of crossed modules (over algebroids) .

In the end of this chapter we give some examples on this notion .

In chapter III , we define the notion of a special double R-algebroid (this is a double R-algebroid with $D_1 = D_2$) and we define a "thin" structure on D which is a morphism

$\theta : \square D_1 \dashrightarrow D$ (where $\square D_1$ is a double R-algebroid with

commutating squares) . An element $\theta(a \begin{smallmatrix} c \\ b \end{smallmatrix} d)$ is called thin ,

where $a, b, c, d \in D_1$. Also we define a connection on D to be a pair of functions $\Gamma, \Gamma' : D_2 \dashrightarrow D$ which satisfy

$$i) \Gamma'a *_2 \Gamma a = c_1 a \quad , \quad \Gamma'a *_1 \Gamma a = c_2 a$$

$$ii) \Gamma'(ab) = (\Gamma'a *_1 c_1 a) *_2 (c_2 a *_1 \Gamma'b)$$

$$\Gamma(ab) = (\Gamma a *_1 c_2 b) *_2 (c_1 b *_1 \Gamma b)$$

$$iii) \Gamma'(a + a_1) *_2 (\alpha +_1 \beta) *_2 \Gamma(d + d_1) = (\Gamma'a *_2 \alpha *_2 \Gamma d) +_2 (\Gamma'a_1 *_2 \beta *_2 \Gamma d_1)$$

where $\alpha, \beta \in D$ with boundaries $(a \begin{smallmatrix} c \\ b \end{smallmatrix} d)$, $(a_1 \begin{smallmatrix} c \\ b \end{smallmatrix} d_1)$ respectively

$$iv) \Gamma'ra *_2 (r \cdot_1 \alpha) *_2 \Gamma rd = r \cdot_2 (\Gamma'a *_2 \alpha *_2 \Gamma d) = \Gamma'a *_2 (r \cdot_2 \alpha) *_2 \Gamma d .$$

Theorem 1 : Let D be a special double R-algebroid with connection Γ' , Γ . Then there is a morphism of special double R-algebroids $\theta : \square D_1 \dashrightarrow D$ which is the identity on D_1 and

$$\Gamma a = \theta(a \begin{smallmatrix} a \\ 1 \end{smallmatrix} 1) \quad , \quad \Gamma'b = \theta(1 \begin{smallmatrix} 1 \\ b \end{smallmatrix} b) \quad , \quad \text{where } a, b \in D_1 .$$

In fact , the reason for defining these two structures on D is that ; the axioms for connection do not involve the addition or scalar multiplication , where as the axioms for a thin structure do ; also it is easier to deal with connection than with thin structure .

Also we define a morphism between two special double R -algebroids with connection . Thus we have a category \underline{DA} of special double R -algebroids with connection . Then we get a functor from the category of special double R -algebroids with connection to the category of crossed modules .

Now , to get a functor from the category of crossed modules to the category of special double R -algebroids with connection we introduce the notion of a "folding operation: Φ which has the effect of "folding" all edges $\alpha \in D$ onto the edge $\partial_1^0 \alpha$. We prove ;

Proposition: There exists a functor from the category of crossed modules to the category of special double R -algebroids with connections .

In fact , we prove

Theorem 2: These functors are equivalent .

Finally we introduce the notion of a reflection on a special double R -algebroid which gives an equivalence between the two algebroid structures .

In chapter IV , we define an ω -algebroid (without connections) by using the cubical complex idea namely ;

An ω -algebroid (without connections) $\underline{A} = \{A_n; \partial_i^\alpha, \epsilon_i\}$ is a cubical complex and for $n \geq 1$, A_n has n algebroid structures over A_{n-1} of the form $(A_n, +_i, *_i, \cdot_i, \partial_i^0, \partial_i^1, \epsilon_i)$ related appropriately to each other and to $\partial_i^0, \partial_i^1, \epsilon_i$.

Thus we can define finite dimensional versions of the above definition . Therefore we get ;

Algebras \rightarrow Algebroids \rightarrow Double algebroids \rightarrow ω -Algebroids .

Also we define a crossed complex \underline{M} (over algebroid) to consist of a sequence of morphisms of R-algebroids over M_0

$$\underline{M}: \dots \rightarrow M_n \xrightarrow{\mathcal{S}} M_{n-1} \xrightarrow{\mathcal{S}} \dots \xrightarrow{\mathcal{S}} M_2 \xrightarrow{\mathcal{S}} M_1$$

satisfying the relations ;

i) each $\mathcal{S} : M_n \rightarrow M_{n-1}$, $n \geq 2$ is the identity on M_0 .

ii) M_1 operates on the right and on the left on each M_n

($n \geq 2$) , by actions $(a, m) \rightarrow a_m$, $(m, b) \rightarrow m^b$,

whenever $m \in M_n(x, y)$, $a \in M_1(w, x)$, $b \in M_1(y, z)$.

iii) If $m \in M_n(x, y)$, $m' \in M_2(y, z)$, $m'' \in M_2(w, x)$, then

$${}_m \mathcal{S} m' = \begin{cases} 0_{xz} & \text{if } n \geq 3 \\ m m' & \text{if } n=2 \end{cases}$$

$$\mathcal{S} m''_m = \begin{cases} 0_{wy} & \text{if } n \geq 3 \\ m'' m & \text{if } n=2 \end{cases}$$

Finally we prove that ;

Theorem 3: There exists a functor γ from the category of ω -algebroids (without connections) to the category of crossed complexes (over algebroids) .

In chapter V , §1 we define an ω -algebroid with connections and the morphisms between them and also we give the definition of a finite dimensional versions of an ω -algebroid .

In § 2 we introduce the notion of "folding operation" Φ , which has a similar effect to the folding operation in the two dimensional case . Also we give the relations between this operation and the axioms of 3 and 4 - tuple algebroids , that is ,

Proposition: Let $a \in A_n$. Then Φa belongs to the associated crossed complex γA .

Proposition: i) If $a, b \in A_n$ with $\partial_j^\alpha a = \partial_j^\alpha b$, for $\alpha = 0, 1$, then $\Phi(a +_j b) = \Phi a +_n \Phi b$.

ii) For $n = 3, 4$, if $a, b \in A_n$ with $\partial_j^1 a = \partial_j^0 b$, then

$$\Phi(a *_j b) = u_j^a (\Phi b) +_n (\Phi a) v_j^b$$

where $u_j^a = \partial_1^0 \dots \partial_{j-1}^0 \partial_{j+1}^0 \dots \partial_n^0 a$ and $v_j^b = \partial_1^1 \dots \partial_{j-1}^1 \partial_{j+1}^1 \dots \partial_n^1 b$.

iii) If $a \in A_n$ and $r \in R$, then

$$\Phi(r \cdot_j a) = r \cdot_n \Phi a.$$

Proposition: 1) For $n = 3$, let $a \in A_2$. Then

$\Phi \epsilon_i a = \Phi \Gamma_j a = \Phi \Gamma'_j a = 0$ in dimension 3 for $1 \leq i \leq 3$ and $1 \leq j \leq 2$.

2) For $n = 4$, let $a \in A_3$. Then

$\Phi \epsilon_i a = \Phi \Gamma_j a = \Phi \Gamma'_j a = 0$ in dimension 4 for $1 \leq i \leq 4$ and $1 \leq j \leq 3$.

Also we define a thin structure on A as follows ;

let $a \in A_n$, then a is called thin if and only if $\Phi a = 0$.

In §3 we construct the coskeleton in terms of "shells" for an n -tuple algebroid and we define ∂_i^α , ϵ_i , Γ_i , Γ'_i and the operations on $\square A_n$ to prove the following ;

Proposition: If (A_n, \dots, A_0) is an n -tuple algebroid, then $(\square A_n, A_n, \dots, A_0)$ is an $(n+1)$ -tuple algebroid.

Proposition: Let A be an ω -algebroid and let $M = \gamma A$ be its associated crossed complex. Let $a \in \square A_{n-1}$ and $\xi \in M_n(u, v)$ where $u = \beta_0 a$, $v = \beta_1 a$. Then a necessary and sufficient

condition for the existence of $b \in A_n$ such that $\partial b = a$ and $\Phi b = \xi$ is that $\xi\xi = \xi\Phi\partial a$. Further if b exist, it is unique.

In §4 we construct a functor λ from the category of 3-truncated crossed complexes to the category of 3-tuple algebroids by using the folding operation. Also we prove that ;

Theorem 4: The functors γ , λ form an adjoint equivalence

$$\gamma : (\underline{\omega}\text{-Alg})^3 \longleftrightarrow (\underline{\text{Crs}})^3 : \lambda .$$