

On Yetter's Invariant and an Extension of the Dijkgraaf-Witten Invariant to Categorical Groups

João Faria Martins*

*Departamento de Matemática, Instituto Superior Técnico,
Av. Rovisco Pais, 1049-001 Lisboa, Portugal.
jmartins@math.ist.utl.pt*

Timothy Porter

*Department of Mathematics, University of Wales, Bangor,
Dean St., Bangor, Gwynedd LL57 1UT, UK.
t.porter@bangor.ac.uk*

August 22, 2006

Abstract

We give an interpretation of Yetter's Invariant of manifolds M in terms of the homotopy type of the function space $\text{TOP}(M, B(\mathcal{G}))$, where \mathcal{G} is a crossed module and $B(\mathcal{G})$ is its classifying space. We use this interpretation to define a twisting of Yetter's Invariant by cohomology classes of crossed modules, defined as cohomology classes of their classifying spaces, in the form of a state sum invariant. In particular, we obtain an extension of the Dijkgraaf-Witten Invariant of manifolds to categorical groups. The straightforward extension to crossed complexes is also considered.

2000 Mathematics Subject Classification: 18F99; 57M27; 57R56; 81T25.

*Financed by Fundação para a Ciência e Tecnologia (Portugal), post-doctoral grant number SFRH/BPD/17552/2004, part of the research project POCTI/MAT/60352/2004 ("Quantum Topology"), also financed by FCT.

Introduction

Let G be a finite group. In the context of Topological Gauge Field Theory, R. Dijkgraaf and E. Witten defined in [DW] a 3-dimensional manifold invariant for each 3-dimensional cohomology class of G . When M is a triangulated manifold, the Dijkgraaf-Witten Invariant can be expressed in terms of a state sum model, reminiscent of Lattice Gauge Field Theory. State sum invariants of manifolds were popularised by V.G. Turaev and O.Ya. Viro, because of their construction of a non-trivial closed 3-manifold state sum invariant from quantum $6j$ -symbols (see [TV]), regularising the divergences of the celebrated Ponzano-Regge Model (see [PR]), which is powerful enough to distinguish homotopic, though non homeomorphic, manifolds. For categorically inclined introductions to state sum invariants of manifolds we refer the reader to [T, BW, Mk1].

State sum models, also known as spin foam models, appear frequently in the context of 3- and 4-dimensional Quantum Gravity, as well as Chern-Simons Theory, making them particularly interesting. See for example [B, Bae] for reviews.

Let us recall the construction of the Dijkgraaf-Witten Invariant. Let G be a finite group. A 3-dimensional group cocycle is a map $\omega: G^3 \rightarrow U(1)$ verifying: $\omega(X, Y, Z)\omega(XY, Z, W)^{-1}\omega(X, YZ, W)\omega(X, Y, ZW)^{-1}\omega(Y, Z, W) = 1$, for any $X, Y, Z, W \in G$. These group 3-cocycles can be seen as 3-cocycles of the classifying space $B(G)$ of G , by considering the usual simplicial structure on $B(G)$. See for example [We], Example 8.1.7.

Let M be an oriented closed manifold. Choose a triangulation of M , and consider a base point $*$ for M so that $*$ is a vertex of M . Suppose that we are provided with a total order on the set of vertices of M . A G -colouring of M is given by an assignment of an element of G to each edge of M , satisfying the flatness condition shown in figure 1. A G -colouring of M therefore defines a morphism $\pi_1(M, *) \rightarrow G$. Let $T = (abcd)$, where $a < b < c < d$, be a tetrahedron of M . The colouring of each edge of T is determined by the colouring of the edges (ab) , (bc) and (dc) , of T . See figure 2. Let $\omega: G^3 \rightarrow U(1)$ be a 3-dimensional group cocycle, representing some 3-dimensional cohomology class of G . The “topological action” $S(T, \omega)$ on a tetrahedron T is $\omega(X, Y, Z)^{\pm 1}$, where $X, Y, Z \in G$ are as in figure 2. Here the exponent on $\omega(X, Y, Z)$ is 1 or -1 according to whether the orientation on T induced by the total order on the set of vertices of M coincides or not with the orientation of M . The expression of the Dijkgraaf-Witten Invariant




Figure 1: Flatness condition imposed on G -colourings of a triangulated manifold M . Here $X, Y \in G$ and $a < b < c$ are vertices of M .




Figure 2: Edges of a tetrahedron of M whose colours determine the colouring of all other edges. Here $a < b < c < d$ are vertices of M .

is the state sum:

$$DW(M, \omega) = \frac{1}{\#G^{n_0}} \sum_{G\text{-colourings}} \prod_{\text{tetrahedra } T} S(T, \omega),$$

where n_0 is the number of vertices of the chosen triangulation of M . This state sum is triangulation independent.

A homotopy theoretic expression for the Dijkgraaf-Witten Invariant is:

$$DW(M) = \sum_{f \in [(M, *), (B(G), *)]} \langle o_M, f^*(w) \rangle,$$

where $o_M \in H_3(M)$ denotes the orientation class of M . The notation $[(M, *), (B(G), *)] = \pi_0(\text{TOP}((M, *), (B(G), *)))$ stands for the set of based homotopy classes of maps $(M, *) \rightarrow (B(G), *)$, where, as before, $B(G)$ denotes the classifying space of G . The equivalence between the two formulations is a consequence of the fact that there exists a one-to-one correspondence between group morphisms $\pi_1(M, *) \rightarrow G$ and pointed homotopy classes of maps $(M, *) \rightarrow (B(G), *)$.

In [Y1], D.N. Yetter defined a 3-dimensional TQFT, which included the $\omega = 1$ case of the Dijkgraaf-Witten Invariant, though he did not consider group cohomology classes. The same kind of construction appeared upgraded in [Y2], where D.N. Yetter extended his framework to handle categorical groups. Recall that categorical groups are equivalent to crossed modules.

Let M be a piecewise linear compact manifold equipped with a triangulation. The definition of Yetter's Invariant (which we will formulate carefully in this article) is analogous to the definition of the Dijkgraaf-Witten Invariant for the case $\omega = 1$. However, in Yetter's construction, in addition to colourings of the edges of M , now by objects of a categorical group, we also have colourings on the faces the triangulation of M by morphism of it, and the flatness condition is transported to the tetrahedrons of M .

The meaning of Yetter's construction was elucidated by the second author in [P1, P2], where an extension Yetter's Invariant to handle general models of n -types appears. The construction of M. Mackaay in [Mk2] is related, conjecturally, with the $n = 3$ case of the construction in [P2], with a further twisting by cohomology classes of 3-types, in the Dijkgraaf-Witten way.

In [FM1, FM2], D.N. Yetter's ideas were applied to 2-dimensional knot theory, and yielded interesting invariants of knotted embedded surfaces in S^4 , with nice calculational properties. In [FM2, FM3] the general formulation Yetter's Invariant for CW-complexes and crossed complexes appeared.

One of the open problems posed by D.N. Yetter at the end of [Y2] was whether his construction could, similarly to the Dijkgraaf-Witten Invariant, be twisted by cohomology classes, and he asked for the right setting for doing this. In this article we formulate such a twisting of Yetter's Invariant, thereby extending the Dijkgraaf-Witten Invariant of 3-manifolds to categorical groups. Note that the (co)homology of a crossed module is defined as being the (co)homology of its classifying space. See for example [E, Pa].

In the course of this work, we prove a new homotopy theoretical interpretation of Yetter's Invariant (Theorem 18), therefore giving a new proof of its existence. See [P1, P2] for alternative interpretations. Our interpretation of Yetter's Invariant extends in the obvious way to the homologically twisted case (Theorem 29), and has close similarities with the interpretation of the Dijkgraaf-Witten Invariant shown above. The main difference is that it is set in the unbased case, and depends on the homotopy groups of each connected component of the function space $\text{TOP}(M, B(\mathcal{G}))$. Here $B(\mathcal{G})$ is the classifying space of the crossed module \mathcal{G} . Our construction does not depend on the dimension of the manifolds, and can also be adapted to the general crossed complex context, which we expect to be closely related with the invariants appearing in [P2, Mk2]. This extension appears in this article.

As we mentioned before, the category of categorical groups is equivalent to the category of crossed modules, which are particular cases of crossed complexes. Crossed complexes \mathcal{A} admit classifying spaces $B(\mathcal{A})$, which generalise

Eilenberg-Mac Lane spaces. The classifying spaces of crossed complexes \mathcal{A} were extensively studied by R. Brown and P.J. Higgins. One of their main results is the total description in algebraic terms of the weak homotopy type of function space $\text{TOP}(M, B(\mathcal{A}))$, where M is a CW-complex. This result together with the General van Kampen Theorem (also due to R. Brown and P.J. Higgins) are the main tools that we will use in this article.

We will make our exposition as self contained as possible, giving throughout description of the concepts of the theory of crossed complexes that we will need, constructions which are due mainly to R. Brown and P.J. Higgins.

This article should be compared to [P3, PT], where similar ideas are used to define Formal Homotopy Quantum Field Theories with background a 2-type.

Contents

1	Crossed Modules and Yetter's Invariant	6
1.1	Crossed Modules and Crossed Complexes	6
1.1.1	Definition of Crossed Modules and Crossed Complexes	6
1.1.2	The General van Kampen Theorem	8
1.1.3	Homotopy of Maps from Crossed Complexes into Crossed Modules	8
1.2	Yetter's Invariant of Manifolds	10
1.2.1	Reduced Crossed Modules and Categorical Groups . .	10
1.2.2	\mathcal{G} -colouring and Yetter's Invariant	11
1.3	Classifying Spaces of Crossed Modules	14
1.3.1	The Homotopy Addition Lemma	14
1.3.2	The Definition of $B(\mathcal{G})$ and Some Properties	15
1.3.3	The Fundamental Groups of Classifying Spaces of Crossed Modules	18
1.3.4	An Interpretation of Yetter's Invariant	19
1.3.5	An Extension of Yetter's Invariant to Crossed Complexes	22
2	Homological Twisting of Yetter's Invariant	23
2.1	Low Dimensional Cohomology of Crossed Modules	23
2.1.1	Homology of Simplicial Sets	24
2.1.2	Homology of Crossed Modules	24
2.1.3	Explicit Description of the Low Dimensional Coboundary Maps	31
2.2	A Homotopy Invariant of 3-Manifolds	32

1 Crossed Modules and Yetter's Invariant

1.1 Crossed Modules and Crossed Complexes

1.1.1 Definition of Crossed Modules and Crossed Complexes

Let G be a groupoid with object set P . We denote the source and target maps of G by $s, t: G \rightarrow P$, respectively. If a groupoid E is totally disconnected, we denote both maps (which are by definition equal) by $\beta: E \rightarrow P$.

Let G and E be groupoids, over the same set P , with E totally disconnected. A crossed module $\mathcal{G} = (G, E, \partial, \triangleright)$ is given by an object preserving groupoid morphism $\partial: E \rightarrow G$ and a left groupoid action \triangleright of G on E by automorphisms. The conditions on \triangleright and ∂ are:

1. $\partial(X \triangleright e) = X\partial(e)X^{-1}$, if $X \in G$ and $e \in E$ are such that $t(X) = \beta(e)$,
2. $\partial(e) \triangleright f = efe^{-1}$, if $e, f \in E$ verify $\beta(e) = \beta(f)$.

See [BI], section 1. Notice that for any $e \in E$ we must have $s(\partial(e)) = t(\partial(e)) = \beta(e)$.

A crossed module is called reduced if P is a singleton. A morphism $F = (\phi, \psi)$ between the crossed modules $\mathcal{G} = (G, E, \partial, \triangleright)$ and $\mathcal{G}' = (G', E', \partial', \triangleright')$ is given by a pair of groupoid morphisms $\phi: G \rightarrow G'$ and $\psi: E \rightarrow E'$, making the diagram:

$$\begin{array}{ccc} E & \xrightarrow{\psi} & E' \\ \partial \downarrow & & \downarrow \partial' \\ G & \xrightarrow{\phi} & G' \end{array}$$

commutative. In addition we must have:

$$t(X) = \beta(e) \implies \phi(X) \triangleright' \psi(e) = \psi(X \triangleright e), \forall X \in G, \forall e \in E.$$

There exists an extensive literature on (reduced) crossed modules. We refer for example to [Ba1, Ba2, BHu, FM2]. The non-reduced case, important for this article, is considered in [B1, B4, BH1, BH2, BHS, BI].

A natural generalisation of the concept of a crossed module is a crossed complex.

Definition 1 A crossed complex \mathcal{A} is given by a complex of groupoids over the same set A_0 (in other words A_0 is the object set of all groupoids), say:

$$\dots \xrightarrow{\partial_{n+1}} A_n \xrightarrow{\partial_n} A_{n-1} \xrightarrow{\partial_{n-1}} A_{n-2} \xrightarrow{\partial_{n-2}} \dots \xrightarrow{\partial_3} A_2 \xrightarrow{\partial_2} A_1,$$

where all boundary maps $\partial_n, n \in \mathbb{N}$ are object preserving, and the groupoids $A_n, n \geq 2$ are totally disconnected. The remaining conditions are:

1. For each $n > 1$, there exists a left groupoid action $\triangleright = \triangleright_n$ of the groupoid A_1 on A_n , by automorphisms, and all the boundary maps $\partial_n, n = 2, 3, \dots$ are A_1 -module morphisms.
2. The groupoid map $A_2 \xrightarrow{\partial_2} A_1$ together with the action \triangleright of A_1 in A_2 defines a crossed module.
3. The groupoid A_n is abelian if $n > 2$.
4. $\partial_2(A_2)$ acts trivially on A_n for $n > 2$.

For a positive L , an L -truncated crossed complex is a crossed complex such that $A_n = \{0\}$ if $n > L$. In particular crossed modules are equivalent to 2-truncated crossed complexes. A crossed complex is called reduced if A_0 is a singleton.

A natural example of a crossed complex is the following one, introduced by A.L. Blakers in [B1], in the reduced case, and extensively studied by J.H.C. Whitehead in [W1, W2], and H.J. Baues, see [Ba1, Ba2]. The non-reduced case, important for this work, is treated in [B1, B4, BH2, BH3, BH4, BH5].

Example 2 Let M be a locally path connected space, so that each connected component of M is path connected. A filtration of M is an increasing sequence $\{M_k, k = 0, 1, 2, \dots\}$ of locally path-connected subspaces of M , whose union is M . In addition, we suppose that M_0 has a non-empty intersection with any connected component of M_k , for each $k \in \mathbb{N}$. Then the sequence of groupoids $\pi_n(M_n, M_{n-1}, M_0)$, with the obvious boundary maps, and natural left actions of the fundamental groupoid $\pi_1(M_1, M_0)$ on them, is a crossed complex, which we denote by $\Pi(M)$, and call the “Fundamental Crossed Complex of the Filtered Space M ”. Note that the assignment $M \mapsto \Pi(M)$, is functorial.

Remark 3 *Let M be a CW-complex. The notation $\Pi(M)$ will always mean the fundamental crossed complex of the skeletal filtration $\{M^k, k \in \mathbb{N}\}$ of M , for which M^k is the k -skeleton of M . Note that it makes sense to consider the fundamental crossed complex of this type of filtrations.*

It is easy to show that crossed complexes and their morphisms, defined in the obvious way, form a category. We will usually denote a crossed complex \mathcal{A} by $\mathcal{A} = (A_n, \partial_n, \triangleright_n)$, or more simply by (A_n, ∂_n) , or even (A_n) . A morphism $f: \mathcal{A} \rightarrow \mathcal{B}$ of crossed complexes will normally be denoted by $f = (f_n)$.

Let $\mathcal{A} = (A_n, \partial_n, \triangleright_n)$ be a crossed complex. For any $a \in A_0$ and any $n \geq 1$, denote $A_n(a)$ as being $\{a_n \in A_n: s(a_n) = a\}$, which is therefore a group if $n \geq 2$. Denote also $A_1(a, b) = \{a_1 \in A_1: s(a_1) = a, t(a_1) = b\}$, where $a, b \in A_0$.

1.1.2 The General van Kampen Theorem

The category of crossed complexes is a category with colimits. See [BH2], section 6. Under mild conditions, the functor Π from the category of filtered spaces to the category of crossed complexes preserves colimits. This fact is known as the General van Kampen theorem, and is due to R. Brown and P.J. Higgins. See [B1, B3, B4, BH1, BH2, BH3, BH4], and also [Ba1, Ba2]. A useful form of the General van Kampen Theorem is the following:

Theorem 4 (R. Brown and P.J. Higgins) *Let M be a CW-complex. Let also $\{M_\lambda, \lambda \in \Lambda\}$ be a covering of M by subcomplexes of M . Then the following diagram:*

$$\bigsqcup_{(\lambda_1, \lambda_2) \in \Lambda^2} \Pi(M_{\lambda_1} \cap M_{\lambda_2}) \rightrightarrows \bigsqcup_{\lambda \in \Lambda} \Pi(M_\lambda) \longrightarrow \Pi(M),$$

with arrows induced by inclusions, is a coequaliser diagram in the category of crossed complexes.

This is corollary 5.2 of [BH2].

1.1.3 Homotopy of Maps from Crossed Complexes into Crossed Modules

The main references now are [BI, BH4]. Note that our conventions are slightly different. See also [B1, B4, W1]. Let G and G' be groupoids. Let also E' be a

totally disconnected groupoid on which G' has a left action. Let $\phi: G \rightarrow G'$ be a groupoid morphism. A map $d: G \rightarrow E'$ is said to be a ϕ -derivation if:

1. $\beta(d(X)) = t(\phi(X)), \forall X \in G$.
2. $d(XY) = (\phi(Y)^{-1} \triangleright d(X))d(Y)$ if $X, Y \in G$ are such that $t(X) = s(Y)$.

Definition 5 Let $\mathcal{A} = (A_n)$ be a crossed complex and $\mathcal{G} = (G, E, \partial, \triangleright)$ be a crossed module. Let also $f: \mathcal{A} \rightarrow \mathcal{G}$ be a morphism of crossed complexes. An f -homotopy H is given by two maps:

$$H_0: A_0 \rightarrow G, \quad H_1: A_1 \rightarrow E,$$

such that:

1. $s(H_0(a)) = f_0(a), \forall a \in A_0$,
2. $H_1: A_1 \rightarrow E$ is an f_1 -derivation, here $f = (f_n)$,

in which case we put $f = s(H)$.

Proposition 6 Under the conditions of the previous definition, suppose that H is an f -homotopy. The map $g = t(H): \mathcal{A} \rightarrow \mathcal{G}$ such that:

1. $g_0(a) = t(H_0(a)), \forall a \in A_0$,
2. $g_1(X) = H_0(s(X))^{-1} f_1(X) (\partial \circ H_1)(X) H_0(t(X)), \forall X \in A_1$,
3. $g_2(e) = H_0(\beta(e))^{-1} \triangleright (f_2(e) (H_1 \circ \partial)(e)), \forall e \in A_2$,

is a morphism of crossed complexes. Moreover, this construction defines a groupoid $\text{CRS}_1(\mathcal{A}, \mathcal{G})$, whose objects are given by all morphisms $\mathcal{A} \rightarrow \mathcal{G}$, and given two morphisms $f, g: \mathcal{A} \rightarrow \mathcal{G}$, a morphism $H: f \rightarrow g$ is an f -homotopy H with $t(H) = g$.

For each $f \in \text{CRS}_0(\mathcal{A}, \mathcal{G})$ (the set of crossed complex morphisms $\mathcal{A} \rightarrow \mathcal{G}$), denote the set of all f -homotopies by $\text{CRS}_1(\mathcal{A}, \mathcal{G})(f)$. Analogously, if $f, g \in \text{CRS}_0(\mathcal{A}, \mathcal{G})$, we denote $\text{CRS}_1(\mathcal{A}, \mathcal{G})(f, g)$ as being the set of elements of the groupoid $\text{CRS}_1(\mathcal{A}, \mathcal{G})$ with source f and target g .

Definition 7 Let $\mathcal{A} = (A_n)$ be a crossed complex and $\mathcal{G} = (G, E, \partial, \triangleright)$ be a crossed module. Let also $f: \mathcal{A} \rightarrow \mathcal{G}$ be a morphism. A 2-fold f -homotopy is a map $I: A_0 \rightarrow E$ such that $\beta(I(a)) = f_0(a), \forall a \in A_0$. As usual $f = (f_n)$.

Denote $\text{CRS}_2(\mathcal{A}, \mathcal{G})(f)$ as being the set of all 2-fold f -homotopies. It is a group with pointwise multiplication as product. Therefore the disjoint union:

$$\text{CRS}_2(\mathcal{A}, \mathcal{G}) \doteq \bigsqcup_{f \in \text{CRS}_0(\mathcal{A}, \mathcal{G})} \text{CRS}_2(\mathcal{A}, \mathcal{G})(f)$$

is a totally disconnected groupoid.

Proposition 8 *Under the conditions of the previous definition, let I be a 2-fold f -homotopy. Then $\partial I = (H_1, H_2)$, where:*

1. $H_0(c) = \partial(I(c)), \forall c \in A_0,$
2. $H_1(X) = f_1(X)^{-1} \triangleright [I(s(X))] I(t(X))^{-1}, \forall X \in A_1,$

is an f -homotopy.

Proposition 9 *We continue to use the notation of the previous definition. Let I be a 2-fold f -homotopy. Let H be a homotopy with source g and target f . Then:*

$$(H \triangleright I)(a) = H_1(a) \triangleright I(a), \forall a \in A_0,$$

is a 2-fold g -homotopy. Moreover, this assignment defines a left groupoid action of $\text{CRS}_1(\mathcal{A}, \mathcal{G})$ on $\text{CRS}_2(\mathcal{A}, \mathcal{G})$ by automorphisms, and together with the boundary map $\partial: \text{CRS}_2(\mathcal{A}, \mathcal{G}) \rightarrow \text{CRS}_1(\mathcal{A}, \mathcal{G})$, already described, defines a crossed module $\text{CRS}(\mathcal{A}, \mathcal{G})$.

This is shown in [BH4] in the general case when \mathcal{G} is a crossed complex.

1.2 Yetter's Invariant of Manifolds

1.2.1 Reduced Crossed Modules and Categorical Groups

Let $\mathcal{G} = (G, E, \partial, \triangleright)$ be a reduced crossed module. Here $\partial: E \rightarrow G$ is a group morphism, and \triangleright is a left action of G on E by automorphisms. We can define a strict monoidal category $\mathcal{C}(\mathcal{G})$ from \mathcal{G} . The set of objects of $\mathcal{C}(\mathcal{G})$ is given by all elements of G . Given an $X \in G$, the set of all morphisms with source X is in bijective correspondence with E , and the target of $e \in E$ is $\partial(e^{-1})X$. In other words a morphism in $\mathcal{C}(\mathcal{G})$ “looks like” $X \xrightarrow{e} \partial(e^{-1})X$. Given $X \in G$ and $e, f \in E$ the composition

$$X \xrightarrow{e} \partial(e^{-1})X \xrightarrow{f} \partial(f^{-1})\partial(e^{-1})X$$

is:

$$X \xrightarrow{ef} \partial(ef)^{-1}X.$$

The tensor product has the form:

$$\begin{array}{ccc} X & Y & XY \\ \downarrow e & \otimes & \downarrow f \\ \partial(e^{-1})X & \partial(f^{-1})Y & \partial(e^{-1})X\partial(f^{-1})Y \end{array} = \begin{array}{c} XY \\ \downarrow (X \triangleright f)e \end{array} .$$

From the definition of a crossed module, it is easy to see that we have indeed defined a strict monoidal category. This construction is an old one. The tensor category $\mathcal{C}(\mathcal{G})$ is a categorical group (see [BMk] and [BHS]). It is well known that the categories of crossed modules and of categorical groups are equivalent (see [BHS, BSp, BMk]). We skip further details on this connection since the point we want to emphasise is that we can construct a strict monoidal category $\mathcal{C}(\mathcal{G})$ from any crossed module \mathcal{G} . This also justifies the title of this article.

1.2.2 \mathcal{G} -colouring and Yetter's Invariant

Let $\mathcal{G} = (G, E, \partial, \triangleright)$ be a finite reduced crossed module. Let M be a triangulated piecewise linear manifold or, more generally, a simplicial complex. We suppose that we have a total order on the set of vertices of M . Each non-degenerate¹ simplex K of M thus inherits a total order on its set of vertices. Therefore, we represent each non-degenerate simplex K of M as $K = (abc\dots)$, where $a < b < c\dots$ are vertices of M . A \mathcal{G} -colouring of M , referred to as a ‘‘Formal \mathcal{G} -Map’’ in [P3, PT], is given by an assignment of an element of G to each non-degenerate edge of M and an element of E to each non-degenerate triangle of M , satisfying the compatibility condition shown in figure 3. A \mathcal{G} -colouring of a non degenerate triangle naturally determines a morphism in $\mathcal{C}(\mathcal{G})$, the monoidal category constructed from \mathcal{G} . See figure 3. There is still an extra condition that \mathcal{G} -colourings must satisfy: Given a coloured non-degenerate 3-simplex, one can construct two morphisms ϕ and ψ in $\mathcal{C}(\mathcal{G})$, see figure 4. Note that we need to use the monoidal structure of $\mathcal{C}(\mathcal{G})$ to define ϕ and ψ . These morphisms are required to be equal.




Figure 3: A \mathcal{G} -colouring of an ordered non-degenerate 2-simplex. Here $X, Y \in G, e \in E$ and $a < b < c$ are vertices of M . To this \mathcal{G} -colouring we associate the morphism $XY \xrightarrow{e} \partial(e)^{-1}XY$, in the category $\mathcal{C}(\mathcal{G})$.




Figure 4: A \mathcal{G} -colouring of an ordered non-degenerate tetrahedron. Here $a < b < c < d$. Note that the compatibility condition $ef = (X \triangleright g)h$ ensures that $\partial(f)^{-1}\partial(e)^{-1}XYZ = \partial(h)^{-1}X\partial(g)^{-1}YZ$.

Let us be specific. Given a non-degenerate simplex K of M , denote the colouring of it by $\mathbf{c}(K)$. Then the conditions that a \mathcal{G} -colouring \mathbf{c} must satisfy are:

$$\partial(\mathbf{c}(abc))^{-1}\mathbf{c}(ab)\mathbf{c}(bc) = \mathbf{c}(ac), \quad (1)$$

whenever (abc) (where $a < b < c$) is a non-degenerate 2-simplex of M , and

$$\mathbf{c}(abc)\mathbf{c}(acd) = (\mathbf{c}(ab) \triangleright \mathbf{c}(bcd))\mathbf{c}(abd), \quad (2)$$

whenever $(abcd)$ (where $a < b < c < d$) is a non-degenerate 3-simplex of M .

Theorem 10 (D.N. Yetter) *Let M be a compact piecewise linear manifold. Consider a triangulation of M with n_0 vertices and n_1 edges. Choose a total order on the set of all vertices of M . Let $\mathcal{G} = (G, E, \partial, \triangleright)$ be a finite reduced crossed module. Then the quantity:*

$$I_{\mathcal{G}}(M) = \frac{\#E^{n_0}}{\#G^{n_0}\#E^{n_1}} \# \{\mathcal{G}\text{-colourings of } M\} \quad (3)$$

is a diffeomorphism invariant of M , thus, in particular, it is triangulation independent, and does not depend on the total order chosen on the set of vertices of M .

This result is due to D.N. Yetter, and it is shown in [Y2, P1, P2]. Similar constructions were considered in [FM1, FM2, FM3], and also [P3, PT]. We will refer to the invariant $I_{\mathcal{G}}$, where \mathcal{G} is a finite crossed module, as ‘‘Yetter’s Invariant’’.

D.N. Yetter’s proof of Theorem 18, relied on Alexander Moves on simplicial complexes, therefore staying in the piecewise linear category. In particular it does not make clear whether $I_{\mathcal{G}}$ is a homotopy invariant. On the other hand Yetter’s argument extends $I_{\mathcal{G}}$ to being a TQFT.

In this article we will give a geometric interpretation of Yetter’s Invariant $I_{\mathcal{G}}$, from which will follow a new proof of its existence. The expected fact that $I_{\mathcal{G}}$ is a homotopy invariant is also implied by it. See [P1, P2] for alternative points of view, with a broad intersection with this one. As a consequence, we also have that $I_{\mathcal{G}}(M)$ is well defined, as a homotopy invariant, for any simplicial complex M .

¹We will consider later general simplicial sets, more general than simplicial complexes, therefore this distinction is important.

In the closed piecewise linear manifold case, we will also consider an additional twisting of $I_{\mathcal{G}}(M)$ by cohomology classes of \mathcal{G} , similar to the Dijkgraaf-Witten Invariant of 3-dimensional manifolds, see [DW], thereby giving a solution to one of the open problems posed by D.N. Yetter in [Y2]. This will be the main aim of this article.

1.3 Classifying Spaces of Crossed Modules

1.3.1 The Homotopy Addition Lemma

Let M be a piecewise linear manifold with a triangulation T , or, more generally, a simplicial complex. Suppose that we have a total order on the set of all vertices of T . Simplicial complexes like this will be called ordered.

Recall that we can define a simplicial set D_T out of any ordered simplicial complex T . The set of n -simplices of D_T is given by all non decreasing sequences (a_0, a_1, \dots, a_n) of vertices of T , such that $\{a_0, a_1, \dots, a_n\}$ is a (non-degenerate) simplex of T . The face and degeneracy maps are defined as:

$$\partial_i(a_0, \dots, a_n) = (a_0, \dots, a_{i-1}, a_{i+1}, \dots, a_n), i = 0, \dots, n,$$

and

$$s_i(a_0, \dots, a_n) = (a_0, \dots, a_i, a_i, \dots, a_n), i = 0, \dots, n,$$

respectively. Notice that the geometric realisation $|D_T|$ of D_T is diffeomorphic with M . There exist several references on simplicial sets. We refer for example, [FP, GZ, Ma, We, KP].

Let $|\Delta(n)|$ be the standard geometric n -simplex. Therefore, it is the geometric realisation of the simplicial set $\Delta(n)$. This simplicial set is obtained from the simplicial complex $(012\dots n)$, the n -simplex, by applying the construction above. Each non-degenerate k -face $s = (a_0 a_1 a_2 \dots a_k)$ of $\Delta(n)$ (thus $a_0 < a_1 < \dots < a_k$) determines, in the obvious way, an element $h(s) \in \pi_k(|\Delta(n)|^k, |\Delta(n)|^{k-1}, a_0)$. Recall that if M is a CW-complex then M^k denotes the k -skeleton of it.

By the General van Kampen Theorem, stated in 1.1.2, the crossed complex $\Pi(|\Delta(n)|)$ is free on the set of non-degenerate faces of $\Delta(n)$. See [B1, B4, W1, BH5]. The definition of free crossed complexes appears for example in [B4]. In particular, if $\mathcal{A} = (A_n)$ is a crossed complex, a morphism $\Pi(|\Delta(n)|) \rightarrow \mathcal{A}$ is specified, uniquely, by an assignment of an element of A_k to each non-degenerate k -face of $\Delta(n)$, with the obvious compatibility relations with the boundary maps. To determine these relations, we need to use

the Homotopy Addition Lemma. This lemma explicitly describes the boundary maps of the crossed complex $\Pi(|\Delta(n)|)$. See [Wh], page 175. Let us be explicit in the low dimensional case. We follow the conventions of [BH5, B1].

Lemma 11 (Homotopy Addition Lemma for $n = 2, 3$) *We have:*

$$\partial(h(0123)) = (h(01) \triangleright h(123))h(013)h(023)^{-1}h(012)^{-1}, \quad (4)$$

$$\partial(h(012)) = h(12)h(23)h(13)^{-1}. \quad (5)$$

These equations are analogues of equations (1) and (2). By the General van Kampen Theorem it follows:

Proposition 12 *Let M be a triangulated piecewise linear manifold with a total order on its set of vertices. Let also \mathcal{G} be a reduced crossed module. There exists a one-to-one correspondence between \mathcal{G} -colourings of M and maps $\Pi(M) \rightarrow \mathcal{G}$, where M has the natural CW-decomposition given by its triangulation. This result still holds for general simplicial complexes.*

This is Proposition 2.1 of [P1]. Note that any map $f: \Pi(M) \rightarrow \mathcal{G}$ yields a \mathcal{G} -colouring \mathbf{c}^f of M , where $\mathbf{c}^f(K) = f(h(K))$. Here K is a non-degenerate simplex of M . To prove this we need to use the Homotopy Addition Lemma.

Different interpretations of \mathcal{G} -colourings appear in [P1, P2, P3, PT].

1.3.2 The Definition of $B(\mathcal{G})$ and Some Properties

We follow [BH5, B1, B4]. Another introduction to this subject appears in [P3]. Let \mathcal{A} be a crossed complex. The nerve $N(\mathcal{A})$ of \mathcal{A} is, by definition, the simplicial set whose set of n -simplices is given by all crossed complex morphisms $\Pi(|\Delta(n)|) \rightarrow \mathcal{A}$, with the obvious face and degeneracy maps. It is a Kan simplicial set. See [BH5]. The classifying space $B(\mathcal{A})$ of \mathcal{A} is the geometric realisation $|N(\mathcal{A})|$ of the nerve of \mathcal{A} . This construction appeared first in [Bl].

Let us unpack this definition in the reduced crossed module case. See [B1], 3.1. Let $\mathcal{G} = (G, E, \partial, \triangleright)$ be a reduced crossed module. The CW-complex $B(\mathcal{G})$ has a unique 0-cell $*$, since $N(\mathcal{G})$ only has one 0-simplex, because \mathcal{G} is reduced. For any $n \in \mathbb{N}$, the set of n -simplices of $N(\mathcal{G})$ is, by Proposition 12, in bijective correspondence with the set of all \mathcal{G} -colourings of $|\Delta(n)|$, and has the obvious faces and degeneracies motivated by this identification.

Consequently, there exists a 1-simplex $s(X)$ of $B(\mathcal{G})$ for any $X \in G$. The 2-simplices of M are given by triples $s(X, Y, e)$, where $X, Y \in G$ and $e \in E$. Here $X = \mathbf{c}(01), Y = \mathbf{c}(12), \partial(e)^{-1}XY = \mathbf{c}(02)$ and $e = \mathbf{c}(012)$, corresponding to the \mathcal{G} -colouring of $|\Delta(2)|$ shown in figure 3. Similarly, the 3-simplex of $N(\mathcal{G})$ determined by the \mathcal{G} -colouring of $|\Delta(3)|$ shown in figure 4 will be denoted by $s(X, Y, Z, e, f, g, h)$, where the relation $ef = (X \triangleright g)h$ must hold. We will describe the set of \mathcal{G} -colourings of the 4-simplex $|\Delta(4)|$ in the following section.

The following result is due to R. Brown and P.J. Higgins. See [BH5], Theorem 2.4, together with its proof.

Theorem 13 *Let C be a simplicial set. In particular the geometric realisation $|C|$ of C is a CW-complex. Let \mathcal{G} be a crossed module. There exists a one-to-one correspondence F between crossed complex morphisms $\Pi(|C|) \rightarrow \mathcal{G}$ and simplicial maps $C \rightarrow N(\mathcal{G})$.*

Remark 14 *In fact, let $g: \Pi(|C|) \rightarrow \mathcal{G}$ be a morphism of crossed complexes. Consider the projection map:*

$$p: \bigsqcup_{n \in \mathbb{N}} C_n \times |\Delta(n)| \rightarrow |C|,$$

according to the usual construction of the geometric realisation of simplicial sets, see for example [Mi1, Ma, FP, GZ]. Note that p is a cellular map, therefore it induces a morphism p_ of crossed complexes. Let $c \in C_n$ be an n -simplex of C , then $F(g)(c): \Pi(|\Delta(n)|) \rightarrow \mathcal{G}$ is the composition:*

$$\Pi(|\Delta(n)|) \xrightarrow{\cong} \Pi(c \times |\Delta(n)|) \xrightarrow{p_*} \Pi(|C|) \xrightarrow{g} \mathcal{G}.$$

These results are a consequence of the version of the General van Kampen Theorem stated in 1.1.2. See [BH5] for details.

Remark 15 *In particular (from Proposition 12), it follows that if M is a piecewise linear manifold with an ordered triangulation T , and C_T is the simplicial set associated with T (see the beginning of 1.3.1), then there exists a one-to-one correspondence between \mathcal{G} -colourings of M and simplicial maps $C_T \rightarrow N(\mathcal{G})$. This result is due to the second author. See [P1, P2].*

The statement of Theorem 13 can be substantially expanded. We follow [BH5]. The assignment $\mathcal{G} \mapsto N(\mathcal{G})$ if \mathcal{G} is a crossed module is functorial.

Similarly, the assignment $C \mapsto \Pi(|C|)$, where C is a simplicial set is also functorial. The second functor is left adjoint to the first one: the Nerve Functor, see [BH5], Theorem 2.4. In fact, the adjunction F of Theorem 13 extends to a simplicial map (which we also call F):

$$N(\text{CRS}(\Pi(|C|), \mathcal{G})) \xrightarrow{F} \text{SIMP}(C, N(\mathcal{G})),$$

which, moreover, is a homotopy equivalence. See [BH5], proof of Corollary 3.5 and of Theorem A. Here if C and D are simplicial sets, then $\text{SIMP}(C, D)$ is the simplicial set whose set of n -simplices is given by all simplicial maps $C \times \Delta(n) \rightarrow D$. See [Ma, GZ], for example. Note that the set of vertices of $N(\text{CRS}(\Pi(|C|), \mathcal{G}))$ is the set $\text{CRS}_0(\Pi(|C|), \mathcal{G})$ of all crossed complex morphisms $\Pi(|C|) \rightarrow \mathcal{G}$. Similarly, there exists a bijective correspondence between vertices of $\text{SIMP}(C, N(\mathcal{G}))$ and simplicial maps $C \rightarrow N(\mathcal{G})$.

The existence of the simplicial homotopy equivalence F is a non trivial fact, due to R. Brown and P.J. Higgins. The proof which appears in [BH5] makes great use of the monoidal closed structure of the category of crossed complexes constructed in [BH4]. A version of the Eilenberg-Zilber Theorem for crossed complexes, representing $\Pi(|C \times D|)$ as a strong deformation retract of $\Pi(|C|) \otimes \Pi(|D|)$, if C and D are simplicial sets, is also required. An approach to this is sketched in [BH5]. A direct proof appears in [To1, To2].

Let C and D be simplicial sets. It is well known that if D is Kan then there exists a weak homotopy equivalence:

$$j: |\text{SIMP}(C, D)| \rightarrow \text{TOP}(|C|, |D|),$$

where the later is given the k -ification of the Compact-Open Topology on the set of all continuous maps $|C| \rightarrow |D|$. See [B2], for example. This weak homotopy equivalence is the composition:

$$|\text{SIMP}(C, D)| \xrightarrow{j_1} |\text{SIMP}(C, S|D)| \xrightarrow{j_2} |S(\text{TOP}(|C|, |D|))| \xrightarrow{j_3} \text{TOP}(|C|, |D|).$$

Here if M is a topological space, then $S(M)$ is the singular complex of M . Since D is a Kan simplicial set, it follows that D is a strong deformation retract of $S(|D|)$, a fact usually known as Milnor's Theorem, see [FP], 4.5. The map j_1 is the geometric realisation of the inclusion $\text{SIMP}(C, D) \rightarrow \text{SIMP}(C, S|D)$, and is therefore a homotopy equivalence.

There exists a one-to-one correspondence between simplicial maps $C \times \Delta(n) \rightarrow S(|D|)$ and continuous maps $|C| \times |\Delta(n)| \cong |C \times \Delta(n)| \rightarrow |D|$. This

fact together with the natural homeomorphism $\text{TOP}(|C| \times |\Delta(n)|, |D|) \rightarrow \text{TOP}(|\Delta(n)|, \text{TOP}(|C|, |D|))$ provides an isomorphism $\text{SIMP}(C, S(|D|)) \rightarrow S(\text{TOP}(|C|, |D|))$, see [GZ], page 132. The map j_2 is the geometric realisation of it, which consequently is a homeomorphism.

The map j_3 is the obvious map $j_3: |S(\text{TOP}(|C|, |D|))| \rightarrow \text{TOP}(|C|, |D|)$. It is well known that the natural map $|S(M)| \rightarrow M$ is a weak homotopy equivalence, for any space M . See [Ma, GZ, FP].

From this discussion, it is obvious that if $f: C \rightarrow D$ is a simplicial map then $j(f) = |f|$. Here $|f|: |C| \rightarrow |D|$ is the geometric realisation of $f: C \rightarrow D$. Therefore:

Theorem 16 *Let C be a simplicial set and \mathcal{G} a reduced crossed module. There exists a weak homotopy equivalence:*

$$\eta: B(\text{CRS}(\Pi(|C|), \mathcal{G})) \rightarrow \text{TOP}(|C|, B(\mathcal{G})),$$

which therefore is a homotopy equivalence when C is finite, because then it follows that $\text{TOP}(|C|, B(\mathcal{G}))$ has the homotopy type of a CW-complex. See [Mi2]. In fact, if we are given a map $g: \Pi(C) \rightarrow \mathcal{G}$, then $\eta(g)$ is the geometric realisation of $F(g): C \rightarrow N(\mathcal{G})$, see Theorem 13.

This is contained in theorem A of [BH5]. Note that $\eta = j \circ |F|$.

1.3.3 The Fundamental Groups of Classifying Spaces of Crossed Modules

Let $\mathcal{G} = (G, E, \partial, \triangleright)$ be a crossed module, not necessarily reduced. Let P be the object set of G and E . There exists a bijective correspondence between 0-cells of $B(\mathcal{G})$ and elements of P .

Theorem 17 *There exists a one-to-one correspondence between connected components of $B(\mathcal{G})$ and connected components of the groupoid G . Given $a \in P$ then:*

$$\pi_1(B(\mathcal{G}), a) = G_{a,a}/\text{Im}\{\partial: E_a \rightarrow G_{a,a}\} \quad (6)$$

and

$$\pi_2(B(\mathcal{G}), a) = \ker\{\partial: E_a \rightarrow G_{a,a}\}. \quad (7)$$

Here $G_{a,b} = \{X \in G: s(X) = a, t(X) = b\}$ and $E_a = \{e \in E: \beta(e) = a\}$, where $a, b \in P$. In fact the groupoid $\pi_1(B(\mathcal{G}), P)$ is the quotient of G by the totally disconnected subgroupoid $\partial(E)$, which is normal in G . All the other homotopy groups of $B(\mathcal{G})$ are trivial.

This is shown in [BH5], in the general case of crossed complexes, and extends to classifying spaces of crossed complexes in the obvious way. It is a consequence of the fact that the nerve $N(\mathcal{A})$ of \mathcal{A} is a Kan simplicial set if \mathcal{A} is a crossed complex.

An immediate corollary of theorems 16 and 17 is that if C is a simplicial set then there exists a one-to-one correspondence between homotopy classes of crossed complex morphisms $\Pi(|C|) \rightarrow \mathcal{G}$ and homotopy classes of maps $|C| \rightarrow B(\mathcal{G})$. More precisely, $f, g: \Pi(|C|) \rightarrow \mathcal{G}$ are homotopic if and only if $\eta(f), \eta(g): |C| \rightarrow B(\mathcal{G})$ are homotopic. R. Brown and P.J. Higgins proved this for any CW-complex, a statement contained in Theorem A of [BH5].

Recall that an n -type is a connected CW-complex M such that $\pi_i(M) = 0, \forall i > n$. Therefore if $\mathcal{A} = (A_m)$ is a crossed complex such that A_m is trivial for $m > n$ then it follows that the classifying space $B(\mathcal{A})$ of \mathcal{A} is an n -type. We warn the reader that this correspondence does not generate all n -types. However, it is well known that crossed modules faithfully represent all 2-types, a result which is due to S. Mac Lane and J.H.C Whitehead. See [MLW]. See for example [Ba1, BH5, L] for alternative proofs.

1.3.4 An Interpretation of Yetter's Invariant

Let \mathcal{G} be a finite reduced crossed module. Recall that $I_{\mathcal{G}}$ stands for Yetter's Invariant. See 1.2.2. We freely use the results and the notation of 1.1.3. We now prove the first main result of this article.

Theorem 18 *Let M be a triangulated piecewise linear manifold or, more generally, a simplicial complex. As usual, we suppose that M is provided with a total order on its set of vertices. Let $\mathcal{G} = (G, E, \partial, \triangleright)$ be a reduced crossed module. We have:*

$$I_{\mathcal{G}}(M) = \sum_{g \in [M, B(\mathcal{G})]} \frac{\#\pi_2(\text{TOP}(M, B(\mathcal{G})), g)}{\#\pi_1(\text{TOP}(M, B(\mathcal{G})), g)}. \quad (8)$$

Here $[M, B(\mathcal{G})] = \pi_0(\text{TOP}(M, B(\mathcal{G})))$ denotes the set of homotopy classes of maps $M \rightarrow B(\mathcal{G})$. In particular, $I_{\mathcal{G}}(M)$ does not depend on the ordered triangulation of M chosen. Moreover, Yetter's Invariant is therefore a homotopy invariant of topological spaces which can be triangulated.

Proof. The two previous theorems will be used. Recall that any simplicial complex naturally defines a simplicial set. See 1.3.1. We freely use

the notation and the results of 1.1.3. We abbreviate $\text{CRS}_2(\Pi(M), \mathcal{G})(f)$ as $\text{CRS}_2(f)$, and analogously for $\text{CRS}_1(\Pi(M), \mathcal{G})(f, g)$ and $\text{CRS}_1(\Pi(M), \mathcal{G})(f)$.

For an $f \in \text{CRS}_0(\Pi(M), \mathcal{G})$, denote $[f]$ as being the set of elements of $\text{CRS}_0(\Pi(M), \mathcal{G})$ which belong to the same connected component of f in the groupoid $\text{CRS}_1(\Pi(M), \mathcal{G})$. Recall also that if $f, g: \Pi(M) \rightarrow \mathcal{G}$ then $\eta(f), \eta(g): M \rightarrow B(\mathcal{G})$ are homotopic if, and only if, f and g belong to the same connected component in the groupoid $\text{CRS}_1(\Pi(M), \mathcal{G})$.

We have:

$$\begin{aligned}
& \sum_{g \in [M, B\mathcal{G}]} \frac{\#\pi_2(\text{TOP}(M, B(\mathcal{G})), g)}{\#\pi_1(\text{TOP}(M, B(\mathcal{G})), g)} \\
&= \sum_{f \in \text{CRS}_0(\Pi(M), \mathcal{G})} \frac{\#\pi_2(\text{TOP}(M, B(\mathcal{G})), \eta(f))}{\#\pi_1(\text{TOP}(M, B(\mathcal{G})), \eta(f))\#[f]} \\
&= \sum_{f \in \text{CRS}_0(\Pi(M), \mathcal{G})} \frac{\#\pi_2(B(\text{CRS}(\Pi(M), \mathcal{G})), f)}{\#\pi_1(B(\text{CRS}(\Pi(M), \mathcal{G})), f)\#[f]} \\
&= \sum_{f \in \text{CRS}_0(\Pi(M), \mathcal{G})} \frac{\#\partial(\text{CRS}_2(f))\#\ker\{\partial: \text{CRS}_2(f) \rightarrow \text{CRS}_1(f, f)\}}{\#\text{CRS}_1(f, f)\#[f]} \\
&= \sum_{f \in \text{CRS}_0(\Pi(M), \mathcal{G})} \frac{\#\text{CRS}_2(f)}{\#\text{CRS}_1(f, f)\#[f]}. \\
&= \sum_{f \in \text{CRS}_0(\Pi(M), \mathcal{G})} \frac{\#\text{CRS}_2(f)}{\#\text{CRS}_1(f)}.
\end{aligned}$$

The last equality follows from the fact that $\text{CRS}_1(\Pi(M), \mathcal{G})$ is a groupoid. Recall that $\text{CRS}_1(f)$ denotes the set of homotopies with source $f: \Pi(M) \rightarrow \mathcal{G}$. The result follows from the following lemmas.

Lemma 19 *Let $\phi: H \rightarrow G$ be a groupoid morphism, with G a group. Let also E be a group on which G acts on the left. Suppose that H is a free groupoid. Any ϕ -derivation $d: H \rightarrow E$ can be specified, uniquely, by its value on the set of free generators of H . In other words, any map from the set of generators of H into E uniquely extends to a derivation $d: H \rightarrow E$.*

Proof. This is shown in [W1] (Lemma 3) for the case in which H is a group, and is easy to prove. The general case can be proved directly in the same

way. Alternatively, we can reduce it to the group case by using the free group \hat{H} on the set of free generators of H . In particular, any groupoid morphism $\phi: H \rightarrow G$ factorises uniquely as $\phi = \psi \circ p$, where p is the obvious map $p: H \rightarrow \hat{H}$, bijective on generators, and $\psi: \hat{H} \rightarrow G$ is a group morphism. Note that if $d: \hat{H} \rightarrow E$ is a ψ -derivation, then $d \circ p: H \rightarrow E$ is a ϕ -derivation. ■

Lemma 20 *There exist exactly $\#E^{n_1}\#G^{n_0}$ f -homotopies if $f: \Pi(M) \rightarrow \mathcal{G}$ is a crossed complex morphism. Here n_0 and n_1 are the number of vertices and edges of M .*

Proof. Follows from the previous lemma and the fact that $\pi_1(M^1, M^0)$ is the free groupoid on the set of edges of M , with one object for each vertex of M . The fact that the crossed module $\mathcal{G} = (G, E, \partial, \triangleright)$ is reduced is being used. ■

Lemma 21 *There exist exactly $\#E^{n_0}$ 2-fold f -homotopies if $f: \Pi(M) \rightarrow \mathcal{G}$ is a crossed complex morphism.*

Proof. This follows from the fact that the crossed module \mathcal{G} is, by assumption, reduced. ■

Using the last two lemmas it follows that:

$$\begin{aligned} & \sum_{f \in [M, B(\mathcal{G})]} \frac{\#\pi_2(\text{TOP}(M, B(\mathcal{G})), f)}{\#\pi_1(\text{TOP}(M, B(\mathcal{G})), f)} \\ &= \sum_{f \in \text{CRS}_0(\Pi(M), \mathcal{G})} \frac{\#\text{CRS}_2(f)}{\#\text{CRS}_1(f)} \\ &= \sum_{f \in \text{CRS}_0(\Pi(M), \mathcal{G})} \frac{\#E^{n_0}}{\#G^{n_0}\#E^{n_1}} \\ &= \frac{\#E^{n_0}}{\#G^{n_0}\#E^{n_1}} \#\{\mathcal{G}\text{-colourings of } M\}. \end{aligned}$$

The last equation follows from Proposition 12. This finishes the proof of Theorem 18. ■

Yetter's Invariant was extended to CW-complexes in [FM2], where an alternative proof of its existence is given, using a totally different argument to the one just shown. Also in the cellular category, an extension of Yetter's Invariant to crossed complexes appeared in [FM3]. In [FM1, FM2], algorithms

are given for the calculation of Yetter's Invariant of complements of knotted embedded surfaces in S^4 , producing non-trivial invariants of 2knots.

Other similar interpretations of Yetter's Invariant appear in [P1, P2].

1.3.5 An Extension of Yetter's Invariant to Crossed Complexes

We can extend Yetter's Invariant of manifolds to handle crossed complexes. Let us sketch how to do this extension. For details (in the pointed CW-complex category) we refer the reader to [FM3].

As part of the information used to construct this extension, we need to know the $n > 3$ cases of the Homotopy Addition Lemma, stated for $n = 2$ and $n = 3$ in 1.3.1. We continue to follow the conventions of [BH5].

Lemma 22 (Homotopy Addition Lemma for $n > 3$) *We have:*

$$\partial h(012\dots n) = h(01) \triangleright h(\partial_0(012\dots n)) + \sum_{i=1}^n (-1)^i h(\partial_i(012\dots n)).$$

*Recall the notation introduced in 1.3.1. Note that this is an identity in the relative fundamental group $\pi_n(|\Delta(n)|^n, |\Delta(n)|^{n-1}, 0 = *)$.*

Definition 23 *Let M be a triangulated piecewise linear manifold with a total order on its set of vertices. Let $\mathcal{A} = (A_i)$ be a reduced crossed complex. An \mathcal{A} -colouring of M is given by an assignment of an element of A_i to each i -simplex of M , which should satisfy the obvious relations motivated by the Homotopy Addition Lemma.*

As before, by the General van Kampen Theorem, it follows that there is a one-to-one correspondence between \mathcal{A} -colourings of M and crossed complex maps $\Pi(M) \rightarrow \mathcal{A}$. Compare with Proposition 12. Using the same techniques that we used in the crossed module case we can prove:

Theorem 24 *Let M be a triangulated compact piecewise linear manifold provided with a total order on its set of vertices, having n_i (non-degenerate) simplices of dimension i for each $i \in \mathbb{N}$. Let also $\mathcal{A} = (A_n)$ be a reduced crossed complex, such that A_n is the trivial group, except for a finite number of $n \in \mathbb{N}$, for which A_n should be finite. The quantity:*

$$I_{\mathcal{A}}(M) = \#\{\mathcal{A}\text{-colourings of } M\} \prod_{i=1}^{\infty} \left(\prod_{j=0}^{\infty} (\#A_{i+j})^{n_j} \right)^{(-1)^i},$$

does not depend on the ordered triangulation of M chosen and it is a homotopy invariant of M , as a topological space. In fact $I_{\mathcal{A}}(M)$ can be interpreted as:

$$I_{\mathcal{A}}(M) = \sum_{f \in \pi_0(\text{TOP}(M, B(\mathcal{A})))} \prod_{i=1}^{\infty} \#\pi_i(\text{TOP}(M, B(\mathcal{A})), f)^{(-1)^i}.$$

Here $B(\mathcal{A})$ is the classifying space of the crossed complex \mathcal{A} . This result still holds in the general case of simplicial complexes.

The proof of this theorem is analogue to the proof of Theorem 18. An important fact to note is that if \mathcal{A} and M are as in Theorem 18, then the function space $\text{TOP}(M, B(\mathcal{A}))$ has only a finite number of connected components, each with a finite number of non-trivial homotopy groups, all of which are finite. This follows from theorems 16 and 17, or more precisely from their extension to the general case of crossed complexes. A proof of Theorem 24 appears in [FM3], in the case when M is a CW-complex with a unique 0-cell. However, using the framework we exposed here in the crossed module case, the proof is easily extended to the case when M may have any finite number of 0-cells.

This construction should be compared with the construction in [P2] of a manifold invariant for any finite n -type.

2 Homological Twisting of Yetter's Invariant

2.1 Low Dimensional Cohomology of Crossed Modules

The (co)homology of crossed modules is studied, for example, in [E, Pa], by considering the (co)homology of their classifying spaces. We do not wish to develop that further. Rather, we will only consider the (co)homology in low dimensions, focusing on a geometric and calculational approach. Our final aim is to define a twisting of Yetter's Invariant by cohomology classes of finite reduced crossed modules, thereby extending the Dijkgraaf-Witten Invariant of manifolds to categorical groups.

Definition 25 *Let \mathcal{G} be a reduced crossed module. The (co)homology of \mathcal{G} is defined as being the (co)homology of the classifying space $B(\mathcal{G})$ of \mathcal{G} , and similarly in the non-reduced case and for crossed complexes.*

2.1.1 Homology of Simplicial Sets

Let D be a simplicial set. It is well known that the homology groups of its geometric realisation $|D|$ can be combinatorially defined from D itself. Let us clarify this statement.

Consider the complex $C(D) = \{C_n(D), \partial_n\}$ of simplicial chains of D . Here $C_n(D)$ is the free abelian group on the set D_n of n -simplices of D . Furthermore $\partial(c) = \sum_{i=0}^n (-1)^i \partial_i(c)$, if $c \in D_n$. See [Ma], 1.2, and [We], 8.2. Note that the assignment $D \mapsto C(D)$, where D is a simplicial set is functorial. The homology of a simplicial set D is defined as being the homology of the chain complex $C(D)$.

The chain complex $C(D)$ has a subcomplex $C^d(D)$, for which $C_n^d(D)$ is the free \mathbb{Z} -module on the set of degenerate n -simplices of D . A classical result asserts that this chain complex is acyclic, see for example [We], proof of Theorem 8.3.8. Therefore, we may equivalently consider the homology of the normalised simplicial chain complex $C^r(D) \doteq C(D)/C^d(D)$. This chain complex is isomorphic with the cellular chain complex of $|D|$, where $|D|$ is provided with its natural cell decomposition coming from the simplicial structure of D . Recall that the geometric realisation $|D|$ of a simplicial set D is a CW-complex with one n -cell for each non-degenerate n -simplex of D . This proves that the simplicial homology of a simplicial set coincides with the cellular homology of its geometric realisation.

In fact a stronger result holds. Namely, there exists an inclusion map $i: C^r(D) \doteq C(D)/C^d(D) \rightarrow C(D)$ such that $p \circ i = \text{id}$, with $i \circ p$ being homotopic to the identity. Here $p: C(D) \rightarrow C^r(D)$ is the projection map. This is the Normalisation Theorem. See [ML], VIII.6.

In this article, we will consider the unnormalised simplicial chain complex.

Note that if M is an ordered simplicial complex, then the simplicial chain complex of D_M , the simplicial set made from M , coincides with its usual definition. See for example [Mau], 4.3. That book discusses both the normalised and unnormalised cases.

2.1.2 Homology of Crossed Modules

Let \mathcal{G} be a crossed module. The classifying space $B(\mathcal{G})$ of \mathcal{G} is the geometric realisation of the simplicial set $N(\mathcal{G})$, the nerve of \mathcal{G} . Therefore, it is natural to consider the simplicial homology of it.

The simplicial structure of $B(\mathcal{G})$ was described in 1.3.2. Let us unpack

the structure of the simplicial chain complex of $B(\mathcal{G})$ for low dimensions. Suppose that $\mathcal{G} = (G, E, \partial, \triangleright)$ is a reduced crossed module. For any $n \in \mathbb{N}$, the simplicial chain group $C_n(\mathcal{G}) \doteq C_n(N(\mathcal{G}))$ is the free abelian group on the set of all \mathcal{G} -colourings of $|\Delta(n)|$, with the obvious boundary maps. In particular:

1. $C_0(\mathcal{G}) = \mathbb{Z}$,
2. $C_1(\mathcal{G})$ is the free \mathbb{Z} -module on the symbols $s(X)$, $X \in G$,
3. $C_2(\mathcal{G})$ the free \mathbb{Z} -module on the symbols $s(X, Y, e)$, where $X, Y \in G$ and $e \in E$,
4. $C_3(\mathcal{G})$ is the free \mathbb{Z} -module on the symbols $s(X, Y, Z, e, f, g, h)$, where $X, Y, Z \in G$ and $e, f, g, h \in E$ must verify $ef = (X \triangleright g)h$.

Moreover we have:

$$\begin{aligned} \partial(s(X)) &= 0, \forall X \in G, \\ \partial(s(X, Y, e)) &= s(X) + s(Y) - s(\partial(e)^{-1}XY), \forall X, Y \in G, \forall e \in E, \end{aligned}$$

and also:

$$\begin{aligned} \partial(s(X, Y, Z, e, f, g, h)) &= s(Y, Z, g) - s(\partial(e)^{-1}XY, Z, f) \\ &\quad + s(X, \partial(g)^{-1}YZ, h) - s(X, Y, e), \end{aligned}$$

where $X, Y, Z \in G$ and $e, f, g, h \in E$ are such that $ef = (X \triangleright g)h$.

The determination of $C_4(\mathcal{G})$ is a bit more complicated. In figures 5 to 9, we display the most general \mathcal{G} -colouring of the 4-simplex (01234). It depends on the variables $X, Y, Z, W \in G$ and $e, f, g, h, i, j, k, m, n, p \in E$, which must satisfy the conditions shown in figures 5 to 9. Namely: $ef = (X \triangleright g)h$, $gi = (Y \triangleright j)k$, $fm = e^{-1}(XY \triangleright j)en$, $hm = (X \triangleright i)p$ and $en = (X \triangleright k)p$. Note that the last relation follows from all the others. From these relations, it is not difficult to conclude that the colourings of all 1-simplices of figures 5 to 9 are coherent. The associated simplicial 4-chains are denoted by $s(X, Y, Z, W, e, f, g, h, i, j, k, m, n, p)$. The determination of the boundary map $\partial: C_4(\mathcal{G}) \rightarrow C_3(\mathcal{G})$ is an easy task.




Figure 5: The most general \mathcal{G} -colouring of (01234): restriction to (0123).




Figure 6: The most general \mathcal{G} -colouring of (01234): restriction to (1234).




Figure 7: The most general \mathcal{G} -colouring of (01234): restriction to (0234).




Figure 8: The most general \mathcal{G} -colouring of (01234): restriction to (0134).




Figure 9: The most general \mathcal{G} -colouring of (01234): restriction to (0124).

2.1.3 Explicit Description of the Low Dimensional Coboundary Maps

Let $\mathcal{G} = (G, E, \partial, \triangleright)$ be a reduced crossed module. We consider the $U(1)$ -cohomology of \mathcal{G} . This is a very particular case of the construction in [Pa]. There, it was considered the general case of cohomology with coefficients in any $\pi_1(B(\mathcal{G}))$ -module. From the discussion above, the group $C^3(\mathcal{G})$ of 3-cochains of \mathcal{G} is given by all maps $\omega: G^3 \times E^4 \rightarrow U(1)$ which verify:

$$\omega(X, Y, Z, e, f, g, h) \neq 1 \implies ef = (X \triangleright g)h.$$

Analogously, the group $C^4(\mathcal{G})$ of 4-cochains is given by all maps $G^4 \times E^{10} \rightarrow U(1)$ which verify:

$$\omega(X, Y, Z, W, e, f, g, h, i, j, k, m, n, p) \neq 1 \implies \begin{cases} ef = (X \triangleright g)h, \\ gi = (Y \triangleright j)k, \\ fm = e^{-1}(XY \triangleright j)en, \\ hm = (X \triangleright i)p, \\ en = (X \triangleright k)p. \end{cases}$$

Recall that the last relation is a consequence of the others. The group $C^2(\mathcal{G})$ of 2-cochains is simply given by all maps $\omega: G^2 \times E \rightarrow U(1)$. The following results follow trivially.

Proposition 26 *Let $\omega \in C^1(\mathcal{G})$ be a 1-cochain. Then:*

$$d\omega(X, Y, e) = \omega(X) + \omega(Y) - \omega(\partial(e)^{-1}XY), \forall X, Y \in G, \forall e \in E.$$

Proposition 27 *Let $\omega \in C^2(\mathcal{G})$ be a 2-cochain. Then*

$$d\omega(X, Y, Z, e, f, g, h) = \omega(Y, Z, g) - \omega(\partial(e)^{-1}XY, Z, f) + \omega(X, \partial(g)^{-1}YZ, h) - \omega(X, Y, e),$$

whenever $ef = (X \triangleright g)h$ and $1 \in \mathbb{C}$ otherwise.

Proposition 28 *Let $\omega \in C^3(\mathcal{G}, A)$ be a 3-cochain. We have:*

$$d\omega(X, Y, Z, W, e, f, g, h, i, j, k, m, n, p) = \omega(X, Y, Z, e, f, g, h) + \omega(Y, Z, W, g, i, j, k) + \omega(X, \partial(g)^{-1}YZ, W, h, m, i, p) - \omega(\partial(e)^{-1}XY, Z, W, f, m, j, n) - \omega(X, Y, \partial(j)^{-1}ZW, e, n, k, p),$$

whenever $ef = (X \triangleright g)h$, $gi = (Y \triangleright j)k$, $fm = e^{-1}(XY \triangleright j)en$, $hm = (X \triangleright i)p$ and $en = (X \triangleright k)p$, and $1 \in \mathbb{C}$ otherwise.

2.2 A Homotopy Invariant of 3-Manifolds

We will restrict our discussion to the 3-dimensional case. However, it is clear that the results that we obtain will still hold, with the obvious adaptations, for any dimension $n \in \mathbb{N}$, and can be extended to handle crossed complexes as well. The n -dimensional analogues of propositions 26 to 28, as well as their extension to crossed complexes, important for calculational purposes, require, however, more laborious calculations.

Let M be a 3-dimensional oriented triangulated closed piecewise linear manifold. The orientation class $o_M \in H_3(M)$ of M chosen can be specified by an assignment of a total order to each non-degenerate tetrahedron of M . These total orders are defined up to even permutations. They define an orientation on each tetrahedron of M . Therefore it is required that if two non-degenerate tetrahedra share a non-degenerate face then the orientations induced on their common face should be opposite.

As usual, we suppose that we are provided with a total order on the set of all vertices of M . Consequently, each non-degenerate 3-simplex K of M can be uniquely represented as $K = (abcd)$ where $a < b < c < d$. If K is a non-degenerate tetrahedron, we say that $r(K)$ is -1 or 1 according to whether the total order induced on the vertices of K differs from the one determined by the orientation of M by an odd or an even permutation.

The orientation class o_M of M , living in the (normalised or unnormalised) simplicial homology group $H_3(M)$ of M is therefore:

$$o_M = \sum_{\text{3-simplices } (abcd)} r(abcd)(abcd),$$

where the sum is extended to the non-degenerate 3-simplices, only.

Let $\mathcal{G} = (G, E, \partial, \triangleright)$ be a finite reduced crossed module. Choose a 3-dimensional cocycle ω representing some cohomology class in $H^3(\mathcal{G})$. For any \mathcal{G} -colouring \mathbf{c} of M define the $U(1)$ -valued ‘‘action’’:

$$\begin{aligned} \mathcal{S}(\mathbf{c}, \omega) &= \prod_{\text{3-simplices } (abcd)} \omega(\mathbf{c}(ab), \mathbf{c}(bc), \mathbf{c}(cd), \mathbf{c}(abc), \mathbf{c}(acd), \mathbf{c}(bcd), \mathbf{c}(abd))^{r(abcd)}, \end{aligned} \tag{9}$$

where the product is extended to the non-degenerate 3-simplices of M , only.

Recall that the set of 3-simplices of the classifying space $B(\mathcal{G})$ of \mathcal{G} is in one-to-one correspondence with the set of \mathcal{G} -colourings of the standard geometric 3-simplex $|\Delta(3)|$. The group of 3-dimensional simplicial cochains of $B(\mathcal{G})$ is given by all assignments of an element of $U(1)$ to each \mathcal{G} -colouring \mathbf{c} of $|\Delta(3)|$. Given a non-degenerate 3-simplex $(abcd)$ of M , the quantity $\omega(\mathbf{c}(ab), \mathbf{c}(bc), \mathbf{c}(cd), \mathbf{c}(abc), \mathbf{c}(acd), \mathbf{c}(bcd), \mathbf{c}(abd))$ is by definition, and under the identification above, exactly $\omega(\mathbf{c}|_{(abcd)})$, where $\mathbf{c}|_{(abcd)}$ is the restriction of the \mathcal{G} -colouring \mathbf{c} of M to the tetrahedron $(abcd)$.

Theorem 29 *Let M be a 3-dimensional closed oriented triangulated piecewise linear manifold, with a total order on its set of vertices. Let n_0 and n_1 be, respectively, the number of vertices and edges of M . Let also $\mathcal{G} = (G, E, \partial, \triangleright)$ be a finite reduced crossed module, and let $w \in H^3(\mathcal{G})$ be a 3-dimensional cohomology class of \mathcal{G} . The quantity:*

$$I_{\mathcal{G}}(M, \omega) = \frac{\#E^{n_0}}{\#G^{n_0}\#E^{n_1}} \sum_{\mathcal{G}\text{-colourings } \mathbf{c}} \mathcal{S}(\mathbf{c}, \omega)$$

is a homotopy invariant of M , and therefore, in particular, it is independent of the ordered triangulation of M chosen. In fact, let $o_M \in H_3(M)$ be the orientation class of M . We have:

$$I_{\mathcal{G}}(M, \omega) = \sum_{g \in [M, B(\mathcal{G})]} \frac{\#\pi_2(\text{TOP}(M, B(\mathcal{G})), g)}{\#\pi_1(\text{TOP}(M, B(\mathcal{G})), g)} \langle o_M, g^*(\omega) \rangle.$$

Here $[M, B(\mathcal{G})] = \pi_0(\text{TOP}(M, B(\mathcal{G})))$ denotes the set of homotopy classes of maps $M \rightarrow B(\mathcal{G})$.

Note that since $N(\mathcal{G})$ is Kan, the Simplicial Approximation Theorem, guarantees that any map $f: M \rightarrow B(\mathcal{G})$ is homotopic to the geometric realisation of a simplicial map $T_M \rightarrow N(\mathcal{G})$, defined up to simplicial homotopy. Here T_M is the simplicial set defined from the triangulation of M . In particular $f^*(\omega)$ is well defined in the simplicial category for any continuous map $f: M \rightarrow B(\mathcal{G})$. The Simplicial Approximation Theorem (for simplicial sets) is proved for example in [S]. Note also the Normalisation Theorem stated in 2.1.1.

The proof of Theorem 29 is analogous to the proof of Theorem 18. The main lemma which we will use for its proof is the following.

Lemma 30 *Let $f \in \text{CRS}_0(\Pi(M), \mathcal{G})$ be a morphism $\Pi(M) \rightarrow \mathcal{G}$. Therefore, by Proposition 12 we can associate a \mathcal{G} -colouring \mathbf{c}^f of M to it. We have:*

$$\langle o_M, \eta(f)^*(w) \rangle = \mathcal{S}(\mathbf{c}^f, \omega).$$

Note that $\eta(f)$ is the realisation of a simplicial map. In fact it is the geometric realisation of $F(f)$. See theorems 13 and 16.

In particular, from theorems 16 and 17 and subsequent comments, it follows that $\mathcal{S}(\mathbf{c}^f, \omega)$ depends only on the homotopy class of $f: \Pi(M) \rightarrow \mathcal{G}$. This also proves that the action $\mathcal{S}(\mathbf{c}^f, \omega)$ does not depend on the cocycle representing the cohomology class $\omega \in H^3(\mathcal{G})$.

Proof. Recall the notation introduced in 1.3.2. The \mathcal{G} -colouring \mathbf{c}^f of M restricts to a \mathcal{G} -colouring $\mathbf{c}^f|_K$ of K , for each non-degenerate simplex K of M . We have:

$$\begin{aligned} & \langle o_M, \eta(f)^*\omega \rangle \\ &= \langle o_M, F(f)^*\omega \rangle, \text{ by definition (c.f. Theorem 16)} \\ &= \langle F(f)_*o_M, \omega \rangle \\ &= \left\langle \sum_{\text{3-simplices } (abcd)} r(abcd)F(f)(abcd), w \right\rangle \\ &= \prod_{\text{3-simplices } (abcd)} \langle F(f)(abcd), w \rangle^{r(abcd)} \\ &= \prod_{\text{3-simplices } (abcd)} \langle f|_{\Pi(abcd)}, w \rangle^{r(abcd)}, \text{ by Theorem 13 and Remark 14} \\ &= \prod_{\text{3-simplices } (abcd)} \langle \mathbf{c}^f|_{(abcd)}, w \rangle^{r(abcd)}, \text{ by Proposition 12} \\ &= \prod_{\text{3-simplices } (abcd)} \omega(\mathbf{c}^f(ab), \mathbf{c}^f(bc), \mathbf{c}^f(cd), \mathbf{c}^f(abc), \mathbf{c}^f(acd), \mathbf{c}^f(bcd), \mathbf{c}^f(abd))^{r(abcd)} \\ &= \mathcal{S}(\mathbf{c}^f, \omega). \end{aligned}$$

The second to last step follows by definition. Note that the sum and the products are to be extended to the non-degenerate 3-simplices of M , only. ■

We now prove Theorem 29.

Proof. (Theorem 29) We maintain the notation we used in the proof of Theorem 18. We have:

$$\begin{aligned}
& \sum_{g \in [M, B(\mathcal{G})]} \frac{\#\pi_2(\text{TOP}(M, B(\mathcal{G})), g)}{\#\pi_1(\text{TOP}(M, B(\mathcal{G})), g)} \langle o_M, g^* \omega \rangle \\
&= \sum_{f \in \text{CRS}_0(\Pi(M), \mathcal{G})} \frac{\#\pi_2(\text{TOP}(M, B(\mathcal{G})), \eta(f))}{\#\pi_1(\text{TOP}(M, B(\mathcal{G})), \eta(f)) \#[f]} \langle o_M, \eta(f)^* \omega \rangle \\
&= \sum_{f \in \text{CRS}_0(\Pi(M), \mathcal{G})} \frac{\#\pi_2(\text{TOP}(M, B(\mathcal{G})), \eta(f))}{\#\pi_1(\text{TOP}(M, B(\mathcal{G})), \eta(f)) \#[f]} \mathcal{S}(c^f, \omega).
\end{aligned}$$

The same calculation as in the proof of theorem 18 finishes the proof. Note that we are implicitly using the fact that if f and f' belong to the same connected component in the groupoid $\text{CRS}_1(\Pi(M), \mathcal{G})$ then it follows that $\eta(f)$ is homotopic to $\eta(f')$ and $\mathcal{S}(c^f, \omega) = \mathcal{S}(c^{f'}, \omega)$. ■

As we referred to before, this theorem can be extended in the obvious way to closed n -manifolds, with n arbitrary, and cohomology classes of crossed complexes. Compare with Theorem 24. It would be interesting to relate our construction with M. Mackaay's one in [Mk2]. Conjecturally, the last one should be related to the 4-manifold invariant obtained from 4-dimensional cohomology classes of crossed complexes of length 3. Finding the precise link forces the determination of all the relation verified by 4-dimensional crossed complex cocycles, which itself requires elaborate calculations. We will consider these issues in a subsequent publication.

Acknowledgements

The first author would like to thank Ronnie Brown, Gustavo Granja, Roger Picken and Marco Mackaay for useful comments.

References

- [Bae] Baez J.C.: Spin Foam Models, Classical Quantum Gravity 15 (1998), no. 7, 1827–1858.
- [B] Barrett J.W.: State Sum Models and Quantum Gravity, XIIIth International Congress on Mathematical Physics (London, 2000), 259–265, Int. Press, Boston, MA, 2001.

- [BMk] Barrett J.W., Mackaay M.: Categorical Representations of Categorical Groups, math.CT/0407463.
- [BW] Barrett J.W., Westbury B.W.: Invariants of Piecewise-Linear 3-Manifolds, *Trans. Amer. Math. Soc.* 348 (1996), no. 10, 3997–4022.
- [Ba1] Baues H.J.: *Combinatorial Homotopy and 4-Dimensional Complexes*. With a preface by Ronald Brown, *de Gruyter Expositions in Mathematics*, 2. Walter de Gruyter & Co., Berlin, 1991.
- [Ba2] Baues H.J.: *Algebraic Homotopy*, Cambridge Studies in Advanced Mathematics, 15. Cambridge University Press, Cambridge, 1989.
- [Bl] Blakers A.L.: Some Relations Between Homology and Homotopy Groups, *Ann. of Math. (2)* 49, (1948). 428–461.
- [B1] Brown R.: Groupoids and Crossed Objects in Algebraic Topology, *Homology Homotopy Appl.* 1 (1999), 1–78 (electronic).
- [B2] Brown R.: *Topology and Groupoids*, Booksurge LLC, S. Carolina, 2006.
- [B3] Brown R.: On the Second Relative Homotopy Group of an Adjunction Space: an Exposition of a Theorem of J. H. C. Whitehead, *J. London Math. Soc. (2)* 22 (1980), no. 1, 146–152.
- [B4] Brown R.: Crossed complexes and Homotopy Groupoids as non Commutative Tools for Higher Dimensional Local-to-Global Problems, Galois theory, Hopf algebras, and semiabelian categories, 101–130, *Fields Inst. Commun.*, 43, Amer. Math. Soc., Providence, RI, 2004.
- [BH1] Brown R., Higgins P.J.: On the Connection Between the Second Relative Homotopy Groups of some Related Spaces, *Proc. London Math. Soc. (3)* 36 (1978), no. 2, 193–212.
- [BH2] Brown R., Higgins P.J.: Colimit Theorems for Relative Homotopy Groups, *J. Pure Appl. Algebra* 22 (1981), no. 1, 11–41.
- [BH3] Brown R., Higgins P.J.: On the Algebra of Cubes, *J. Pure Appl. Algebra* 21 (1981), no. 3, 233–260.

- [BH4] Brown R., Higgins P.J.: Tensor Products and Homotopies for ω -Groupoids and Crossed Complexes, *J. Pure Appl. Algebra* 47 (1987), no. 1, 1–33.
- [BH5] Brown R., Higgins P.J.: The Classifying Space of a Crossed Complex, *Math. Proc. Cambridge Philos. Soc.* 110 (1991), no. 1, 95–120.
- [BHS] Brown R., Higgins P.J., Sivera R.: Nonabelian Algebraic Topology, part I (preliminary version).
- [BHu] Brown R., Huebschmann J.: Identities Among Relations, *Low-dimensional topology (Bangor, 1979)*, pp. 153–202, *London Math. Soc. Lecture Note Ser.*, 48, Cambridge Univ. Press, Cambridge-New York, 1982.
- [BI] Brown R., İçen İ: Homotopies and Automorphisms of Crossed Modules of Groupoids, *Appl. Categ. Structures* 11 (2003), no. 2, 185–206.
- [BSp] Brown R., Spencer C.B.: G -groupoids, Crossed Modules and the Fundamental Groupoid of a Topological Group. *Nederl. Akad. Wetensch. Proc. Ser. A* 79 *Indag. Math.* 38 (1976), no. 4, 296–302.
- [DW] Dijkgraaf R., Witten E.: Topological Gauge Theories and Group Cohomology, *Comm. Math. Phys.* 129 (1990), no. 2, 393–429.
- [E] Ellis G.: Homology of 2-Types, *J. London Math. Soc.* (2) 46 (1992), no. 1, 1–27.
- [FM1] Faria Martins J.: Categorical Groups, Knots and Knotted Surfaces, to appear in *J. Knot Theory Ramifications*. Preliminary version: math.GT/0502562.
- [FM2] Faria Martins J.: On 2-Dimensional Homotopy Invariants of Complements of Knotted Surfaces, math.GT/0507239.
- [FM3] Faria Martins J.: On the Homotopy Type and the Fundamental Crossed Complex of the Skeletal Filtration of a CW-Complex, math.GT/0605364.
- [FP] Fritsch R., Piccinini R. A.: *Cellular Structures in Topology*, Cambridge Studies in Advanced Mathematics, 19. Cambridge University Press, Cambridge, 1990.

- [GZ] Gabriel P., Zisman M.: Calculus of Fractions and Homotopy Theory. *Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 35* Springer-Verlag New York, Inc., New York 1967.
- [KP] Kamps K.H., Porter T.: *Abstract Homotopy and Simple Homotopy Theory*. World Scientific Publishing Co., Inc., River Edge, NJ, 1997.
- [L] Loday J.L.: Spaces with Finitely Many Non-Trivial Homotopy Groups, *J. Pure Appl. Algebra* 24 (1982), 179–202.
- [ML] Mac Lane S.: *Homology*. Reprint of the 1975 edition. *Classics in Mathematics*. Springer-Verlag, Berlin, 1995.
- [MLW] Mac Lane S., Whitehead J.H.C.: On the 3-Type of a Complex, *Proc. Nat. Acad. Sci. U. S. A.* 36, (1950). 41–48.
- [Mk1] Mackaay M.: Spherical 2-Categories and 4-Manifold Invariants, *Adv. Math.* 143 (1999), no. 2, 288–348.
- [Mk2] Mackaay M.: Finite Groups, Spherical 2-Categories, and 4-Manifold Invariants, *Adv. Math.* 153 (2000), no. 2, 353–390.
- [Mau] Maunder C.R.F.: *Algebraic Topology*, Cambridge University Press, Cambridge-New York, 1980.
- [Ma] May J.P.: *Simplicial Objects in Algebraic Topology*, Van Nostrand Mathematical Studies, No. 11 D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto, Ont.-London 1967.
- [Mi1] Milnor J.: The Geometric Realization of a Semi-Simplicial Complex, *Ann. of Math. (2)* 65 (1957), 357–362.
- [Mi2] Milnor J.: On Spaces Having the Homotopy Type of CW-Complex. *Trans. Amer. Math. Soc.* 90 1959 272–280.
- [PR] Ponzano G., Regge T.: *Semiclassical Limit of Racah Coefficients, Spectroscopic and Group Theoretical Methods in Physics* ed F Bloch et al (Amsterdam: North-Holland) 1968.
- [Pa] Paoli S.: (Co)homology of Crossed Modules with Coefficients in a π_1 -Module, *Homology Homotopy Appl.* 5 (2003), no. 1, 261–296 (electronic).

- [P1] Porter T.: Interpretations of Yetter's Notion of G -Coloring: Simplicial Fibre Bundles and Non-Abelian Cohomology, *J. Knot Theory Ramifications* 5 (1996), no. 5, 687–720.
- [P2] Porter T.: Topological Quantum Field Theories from Homotopy n -Types, *J. London Math. Soc. (2)* 58 (1998), no. 3, 723–732.
- [P3] Porter T.: Formal Homotopy Quantum Field Theories, II : Simplicial Formal Maps, *math.QA/0512034*.
- [PT] Porter T., Turaev V.G.: Formal Homotopy Quantum Field Theories, I: Formal Maps and Crossed C-algebras, *math.QA/0512032*.
- [S] Sanderson B.J.: The Simplicial Extension Theorem, *Math. Proc. Cambridge Philos. Soc.* 77 (1975), 497–498.
- [To1] Tonks A.P.: On the Eilenberg-Zilber Theorem for Crossed Complexes, *J. Pure Appl. Algebra* 179 (2003), no. 1-2, 199–220.
- [To2] Tonks A.P.: Theory and Applications of Crossed Complexes, U.W. Bangor Ph.D. thesis (1994), available at: <http://www.informatics.bangor.ac.uk/public/mathematics/research/ftp/theses/tonks.ps.gz>
- [T] Turaev V.G.: Quantum Invariants of Knots and 3-Manifolds. de Gruyter Studies in Mathematics, 18. Walter de Gruyter & Co., Berlin, 1994.
- [TV] Turaev V.G., Viro O.Ya.: State Sum Invariants of 3-Manifolds and Quantum $6j$ -Symbols. *Topology* 31 (1992), no. 4, 865–902.
- [We] Weibel C.A.: An Introduction to Homological Algebra, Cambridge Studies in Advanced Mathematics, 38. Cambridge University Press, Cambridge, 1994.
- [Wh] Whitehead G.W.: Elements of Homotopy Theory, Graduate Texts in Mathematics, 61. Springer-Verlag, New York-Berlin, 1978.
- [W1] Whitehead J.H.C.: Combinatorial Homotopy II, *Bull. Amer. Math. Soc.* 55, (1949). 453–496.

- [W2] Whitehead J.H.C.: Simple Homotopy Types, Amer. J. Math. 72, (1950). 1–57.
- [Y1] Yetter D.N.: Topological Quantum Field Theories Associated to Finite Groups and Crossed G -Sets, J. Knot Theory Ramifications 1 (1992), no. 1, 1–20.
- [Y2] Yetter D.N.: TQFT's from Homotopy 2-Types, J. Knot Theory Ramifications 2 (1993), no. 1, 113–123.